

ON GRADED 2-ABSORBING SUBMODULES OVER
 Gr -MULTIPLICATION MODULES

Fatemeh Soheilnia and Ahmad Yousefian Darani

Abstract. Let G be a multiplicative group with identity e , R be a G -graded commutative ring and M be a graded R -module. The aim of this article is some investigations of graded 2-absorbing submodules over Gr -multiplication modules. A graded submodule N of R -module M is called graded 2-absorbing if whenever $a, b \in h(R)$ and $m \in h(M)$ with $abm \in N$, then either $ab \in (N :_R M)$ or $am \in N$ or $bm \in N$. We also introduce the concept of graded classical 2-absorbing submodule as a generalization of graded classical prime submodules and show a number of results in this class.

1. Introduction

Throughout this paper all rings are G -graded commutative rings and M is a graded R -module with non-zero identity. Let G be a multiplicative group with identity e . Then R is a G -graded ring, if there exist additive subgroups R_g of R such that $R = \bigoplus_{g \in G} R_g$ and $R_g R_h \subseteq R_{gh}$ for all $g, h \in G$. It is denoted by $G(R)$. The elements of R_g are called *homogeneous* of degree g , where R_g are additive subgroups of R indexed by elements $g \in G$. If $a \in R$, then a can be written uniquely as $\sum_{g \in G} a_g$, where a_g is the component of a in R_g . Also, we write $h(R) = \bigcup_{g \in G} R_g$. Let I be an ideal of $G(R)$. Then I is a *graded ideal* of $G(R)$ if $I = \bigoplus_{g \in G} (I \cap R_g)$. Moreover, R_e is a subring of $G(R)$ and $1 \in R_e$.

Let R be a G -graded ring and M be an R -module. Then M is called a *graded R -module* (G -graded R -module) if there exists a family of subgroups $\{M_g\}_{g \in G}$ of M such that $M = \bigoplus_{g \in G} M_g$ (as abelian groups) and $R_g M_h \subseteq M_{gh}$ for all $g, h \in G$. Here, $R_g M_h$ denotes the additive subgroup of M consisting of all finite sums of elements $r_g m_h$ with $r_g \in R_g$ and $m_h \in M_h$. We write $h(M) = \bigcup_{g \in G} M_g$ and the elements of $h(M)$ are called *homogeneous*. If $M = \bigoplus_{g \in G} M_g$ is a graded R -module, then for

2010 Mathematics Subject Classification: 13A02, 16W50

Keywords and phrases: Graded multiplication modules; graded prime submodules; graded 2-absorbing submodules; graded classical submodules; graded classical 2-absorbing submodules.

all $g \in G$ the subgroup M_g of M is an R_e -module. Let $M = \bigoplus_{g \in G} M_g$ be a graded R -module and N be a submodule of M . Then N is called a *graded submodule* of M if $N = \bigoplus_{g \in G} N_g$ where $N_g = N \cap M_g$ for $g \in G$. In this case, N_g is called the g -component of N , (cf. [10]). Moreover, M/N becomes a graded R -module with g -component $(M/N)_g = (M_g + N)/N$ for $g \in G$. Let R be a G -graded ring and M, N be graded R -modules. Then $(N :_R M) = \{r \in h(R) \mid rM \subseteq N\}$ is a graded ideal of $G(R)$ and for every element $a \in h(R)$ and $m \in h(M)$, IN , aN and Rm are graded submodules of M and Ra is a graded ideal, (cf. [3, 4]). *The annihilator* of an element m of M which is $ann(m) = (0 :_R m) = \{r \in h(R) \mid rm = 0\}$. A graded R -module M is said to be a *cancellation graded module* of $G(R)$ if for all graded ideals I and J of $G(R)$ such that $IM = JM$, it follows that $I = J$. A graded R -module M is *faithful*, if for every element $r \in h(R)$ such that $rM = 0$, it follows that $r = 0$. A graded R -module M is called *graded multiplication (Gr-multiplication) R -module*, if for every graded submodule N of M , there exists a graded ideal I of $G(R)$ such that $N = IM$. In this case, we can easily show that if M is a Gr -multiplication R -module, then $N = (N :_R M)M$ for every graded submodule N of M . Let M be a Gr -multiplication R -module, $N = IM$ and $K = JM$ are graded submodules of M where I and J are graded ideals of $G(R)$. The product of N and K is denoted by NK and is defined by $(IJ)M$. Then the product of N and K is independent of presentations of N and K , by [8, Theorem 4]. Moreover, we can define the product of two elements $m, m' \in h(M)$ — if $Rm = IM$ and $Rm' = JM$, then $mm' = (IM)(JM) = (IJ)M$, (cf. [8, 11]).

A graded ideal I of $G(R)$ is said to be a *graded prime (G-prime)* (resp. *graded weakly prime ideal*), if $I \neq R$ and whenever $a, b \in h(R)$ with $ab \in I$ (resp. $0 \neq ab \in I$), then either $a \in I$ or $b \in I$. The graded radical of I , which is denoted by $Gr(I)$, is the set of all $x \in R$ such that for each $g \in G$ there exists $n_g > 0$ with $x_g^{n_g} \in I$. Note that, if r is a homogeneous element of $G(R)$, then $r \in Gr(I)$ if and only if $r^n \in I$ for some $n \in \mathbb{N}$. It is easy to see that if I is a graded ideal of $G(R)$, then $Gr(I)$ is a graded ideal of $G(R)$, [12, Proposition 2.3]. If I is a graded ideal of $G(R)$, then $Gr(I)$ is the intersection of all G -prime ideals of $G(R)$ containing I , [12, Proposition 2.5]. Furthermore, a graded ideal I of $G(R)$ is said to be a *graded primary (G-primary)* (resp. *graded weakly primary ideal*), if $I \neq R$ and whenever $a, b \in h(R)$ such that $ab \in I$ ($0 \neq ab \in I$), then either $a \in I$ or $b \in Gr(I)$. If $Gr(I)$ is a G -prime ideal, then I is called a *graded P-primary (G-P-primary) ideal*. M is a graded maximal (G -maximal) ideal of G -graded ring R , if $M \neq R$ and there is no graded ideal I of $G(R)$ such that $M \subset I \subset R$.

Graded prime submodules on G -graded commutative rings have been introduced and studied in [3, 4, 8]. Let R be a G -graded ring and M be a graded R -module. A proper graded submodule N of M is called *graded prime submodule* of M , if whenever $a \in h(R)$ and $m \in h(M)$ with $am \in N$, then either $a \in (N :_R M)$ or $m \in N$. In this paper, we generalize the concept of *2-absorbing ideals* in [5] to the concept of *graded 2-absorbing submodules*. Refer that the concept of graded 2-absorbing submodule is a characterization of 2-absorbing submodule which has been explained by A. Yousefian Darani and F. Soheilnia in [15, 16]. Later, K. Al-Zoubi and R. Abu-Dawwas in [1],

extended the concept of *graded 2-absorbing submodules*. They defined that a graded proper submodule N of M is said to be a *graded 2-absorbing submodule*, if whenever $a, b \in h(R)$ and $m \in h(M)$ with $abm \in N$, then either $ab \in (N :_R M)$ or $am \in N$ or $bm \in N$. The concept of graded 2-absorbing ($G(2)$ -absorbing) ideal is defined in [13]. Let R be a G -graded ring and I be a graded ideal of $G(R)$. Then I is said to be a *graded 2-absorbing ($G(2)$ -absorbing) ideal* if $I \neq G(R)$ and whenever $a, b, c \in h(R)$ with $abc \in I$, then either $ab \in I$ or $bc \in I$ or $ac \in I$.

The graded classical prime and primary submodules have been introduced and studied in [2] while the concept of classical prime and classical primary submodule were described by Behboodi in [7], M. Baziar and M. Behboodi in [6]. Let R be a G -graded ring and M be a graded R -module. A proper graded submodule N of M is said to be a *graded classical prime* (resp. *graded classical primary*) *submodule*, if whenever $a, b \in h(R)$ and $m \in h(M)$ with $abm \in N$, then either $am \in N$ or $bm \in N$ (resp., $b^n m \in N$ for some positive integer n). Recently, some researchers in [9] have explained and expanded the concept of classical 2-absorbing submodule over commutative rings. Let M be a R -module over commutative ring R and N be a proper submodule. A submodule N of M is called a classical 2-absorbing submodule, if whenever $a, b, c \in R$ and $m \in M$ such that $abcm \in N$, then either $abm \in N$ or $bcm \in N$ or $acm \in N$. Here we introduce the concept of graded classical 2-absorbing submodules of M over G -graded commutative rings. Let R be a G -graded commutative ring, M be a graded R -module and N be a graded proper submodule of M . We say that N is a *classical graded 2-absorbing submodule*, if whenever $a, b, c \in h(R)$ and $m \in h(M)$ with $abcm \in N$, then either $abm \in N$ or $bcm \in N$ or $acm \in N$. We show more results about graded 2-absorbing submodules which are generalizations of Gr -multiplication R -modules over G -graded commutative rings in Section 2. In Section 3, we show more results on graded classical 2-absorbing submodules that are generalizations of graded prime submodules and graded 2-absorbing submodules of graded R -modules over G -graded rings.

2. Properties of graded 2-absorbing submodules

Let R be a G -graded ring and M be a graded R -module. A graded proper submodule N of M is called graded 2-absorbing submodule, if whenever $a, b \in h(R)$ and $m \in h(M)$ such that $abm \in N$, then either $ab \in (N :_R M)$ or $am \in N$ or $bm \in N$. We say that $N_r = (N :_M r) = \{m \in h(M) \mid mI \subseteq N\}$ is a graded 2-absorbing submodule of M , (cf. [14]). To start with, we show a result on graded 2-absorbing submodules.

THEOREM 2.1. *Let R be a G -graded commutative ring, M be a graded R -module and N be a graded proper submodule of M . Suppose that N is a graded 2-absorbing submodule. Then the following statements hold:*

(i) *For all elements $a, b \in h(R)$ and $m \in h(M)$, $ab \notin (N :_R M)$ implies that $(N :_R abm) = (N :_R am) \cup (N :_R bm)$. Moreover, $(N :_R abm) = (N :_R am)$ or $(N :_R abm) = (N :_R bm)$;*

- (ii) For all elements $a, b \in h(R)$, $ab \notin (N :_R M)$ implies that $(N :_M ab) = (N :_M a)$ or $(N :_M ab) = (N :_M b)$;
- (iii) For every element $a \in h(R)$ and graded submodule K of M , $aK \not\subseteq N$ implies that $(N :_R aK) = (N :_R K)$ or $(N :_R aK) = ((N :_M a) :_R M)$;
- (iv) For every element $a \in h(R)$, every element $m \in h(M)$ and every graded ideal I of $G(R)$, $aIm \subseteq N$ implies that $Ia \subseteq (N :_R M)$ or $am \in N$ or $Im \subseteq N$;
- (v) For every graded ideal I, J of $G(R)$ and every element $m \in h(M)$, $IJm \subseteq N$ implies that $IJ \subseteq (N :_R M)$ or $Im \subseteq N$ or $Jm \subseteq N$;
- (vi) For every element $a, b \in h(R)$ and every graded submodule K of M with $abK \subseteq N$ implies that $ab \in (N :_R M)$ or $aK \subseteq N$ or $bK \subseteq N$;
- (vii) For every element $a \in h(R)$, every graded ideal I of $G(R)$ and graded submodule K of M , $aIK \subseteq N$ implies that $aI \subseteq (N :_R M)$ or $aK \subseteq N$ or $IK \subseteq N$;
- (viii) For every graded ideal I of $G(R)$ and graded submodule K of M , $IK \not\subseteq N$, implies that $(N :_R IK) = (N :_R K)$ or $(N :_R IK) = ((N :_M I) :_R M)$;
- (ix) For every graded ideal I, J of $G(R)$ and graded submodule K of M , $IJK \subseteq N$ implies that $IJ \subseteq (N :_R M)$ or $IK \subseteq N$ or $JK \subseteq N$.

Proof. (i) Assume that $a, b \in h(R)$ and $m \in h(M)$ with $ab \notin (N :_R M)$. Let $x \in (N :_R abm)$ and so $xabm \in N$. Since N is a graded 2-absorbing submodule and $ab \notin (N :_R M)$, we conclude that $xam \in N$ (so $x \in (N :_R am)$) or $xbm \in N$ (so $x \in (N :_R bm)$). Then $(N :_R abm) = (N :_R am) \cup (N :_R bm)$. Clearly, $(N :_R abm) = (N :_R am)$ or $(N :_R abm) = (N :_R bm)$.

(ii) Assume that $a, b \in h(R)$ with $ab \notin (N :_R M)$. Let $m \in (N :_M ab)$ be such that $abm \in N$. Since N is a graded 2-absorbing submodule and $ab \notin (N :_R M)$, we conclude that $am \in N$ (so $m \in (N :_M a)$) or $bm \in N$ (so $m \in (N :_M b)$). Then $(N :_M ab) = (N :_M a)$ or $(N :_M ab) = (N :_M b)$.

(iii) Assume that $a \in h(R)$ and K is a graded submodule of M such that $aK \not\subseteq N$. Suppose that $m \in (N :_R aK)$. Then $amK \subseteq N$ and so $K \subseteq (N :_R am)$. If $am \in (N :_R M)$, then $m \in ((N :_R M) :_M a)$. We can suppose that $am \notin (N :_R M)$. Then it follows from (ii) that $(N :_R am) \subseteq (N :_R m)$ or $(N :_R am) \subseteq (N :_M a)$. If $(N :_R am) \subseteq (N :_R m)$, then $K \subseteq (N :_M a)$ and so $aK \subseteq N$, which is a contradiction. Thus $(N :_R am) \subseteq (N :_M a)$ and hence $K \subseteq (N :_R m)$. Then $Km \subseteq N$ and so $m \in (N :_R K)$. Hence $(N :_R aK) \subseteq (N :_R K) \cup ((N :_M a) :_R M)$. The converse inclusion is obvious. Therefore, $(N :_R aK) = (N :_R K)$ or $(N :_R aK) = ((N :_M a) :_R M)$.

(iv) Assume that for $a \in h(R)$, $m \in h(M)$ and a graded ideal I of $G(R)$, $aIm \subseteq N$, $Ia \not\subseteq (N :_R M)$ and $am \notin N$ hold. There exists $b \in I \cap h(R)$ such that $ab \notin (N :_R M)$. Since $bam \in N$, N is a graded 2-absorbing submodule of M , $ab \notin (N :_R M)$ and $am \notin N$, we get that $bm \in N$. Now suppose that there exists $i \in I$ such that $(b+i)am \in N$. Then $(b+i)a \in (N :_R M)$ or $(b+i)m \in N$. If $(b+i)a \in N$, then $im \in N$. If $(b+i)m \in (N :_R M)$, then $ia \notin (N :_R M)$. Since $iam \in N$ and N is a graded 2-absorbing submodule of M , we obtain that $im \in N$. Therefore $Im \subseteq N$, as required.

(v) Assume that $IJm \subseteq N$ for some graded ideals I, J of $G(R)$ and $m \in h(M)$ with $Im \not\subseteq N$ and $Jm \not\subseteq N$. Then there exist $a \in I \setminus h(R)$ and $b \in I \setminus h(R)$ such that $am \notin N$ and $bm \notin N$. Since $aJm \subseteq N$, $am \notin N$ and $Jm \not\subseteq N$, (iv) implies

$aJ \subseteq (N :_R M)$. Then $(I \setminus (N :_R m))J \subseteq (N :_R M)$. Now, since $Ibm \subseteq N$, $bm \notin N$ and $Im \not\subseteq N$, from (iv) we conclude that $Ib \subseteq (N :_R M)$, thus $I(J \setminus (N :_R m)) \subseteq (N :_R M)$. Hence there exist $i \in I \cap h(R)$ and $j \in J \cap h(R)$ such that $aj \in (N :_R M)$ or $ib \in (N :_R M)$. Since $(a+i) \in I$ and $(b+j) \in J$, we have $(a+i)(b+j)m \in N$. Then $(a+i)(b+j) \in (N :_R M)$ or $(a+i)m \in N$ or $(b+j)m \in N$. Since N is a graded 2-absorbing submodule of M , if $(a+i)(b+j) = ab + aj + ib + ij \in (N :_R M)$, then $ij \in (N :_R M)$. If $(a+i)m \in N$, then $im \notin N$ and thus $i \in I \setminus (N :_R m)$ and so $ij \in (N :_R M)$. Similarly, if $(b+j)m \in N$, then $ij \in (N :_R M)$. Therefore $IJ \subseteq (N :_R M)$.

(vi) Assume that $abK \in N$ for some graded submodule K of M and $a, b \in h(R)$ with $ab \notin (N :_R M)$. Then $K \subseteq (N :_M ab)$. Since $ab \notin (N :_R M)$, it follows from (ii) that $K \subseteq (N :_M ab) = (N :_M a)$ or $K \subseteq (N :_M ab) = (N :_M b)$. Hence either $aK \subseteq N$ or $bK \subseteq N$.

(vii) Assume that $aIK \subseteq N$ for some graded ideal I of $G(R)$, graded submodule K of M and $a \in h(R)$ such that $aI \not\subseteq (N :_R M)$. Then there exists $i \in I$ such that $ai \notin (N :_R M)$. Since $aiK \subseteq N$ and $ai \notin (N :_R M)$, then either $aK \subseteq N$ or $iK \subseteq N$ (so $IK \subseteq N$).

(viii) Assume that I is a graded ideal and K is a graded submodule of M such that $IK \not\subseteq N$. Suppose that $a \in (N :_R IK)$. Then $aIK \subseteq N$ and so (vii) implies $aI \subseteq (N :_R M)$ or $aK \subseteq N$. Thus $aIM \subseteq N$ (so $a \in ((N :_M I) :_R M)$) or $a \in (N :_R K)$. Hence $(N :_R IK) \subseteq (N :_R K) \cup ((N :_M I) :_R M)$. The converse inclusion is obvious, hence $(N :_R IK) = (N :_R K)$ or $(N :_R IK) = ((N :_M I) :_R M)$.

(ix) Assume that $IJK \subseteq N$ for some graded ideals I, J of $G(R)$ and graded submodule K of M with $IJ \not\subseteq (N :_R M)$. Then there exist $a \in I \cap h(R)$ and $b \in J \cap h(R)$ such that neither $aK \in N$ nor $bK \in N$. Since $abK \in N$ and neither $aK \in N$ nor $bK \in N$, from (vi) we get that $ab \in (N :_R M)$. Since $IJ \not\subseteq (N :_R M)$, we obtain that $rs \notin (N :_R M)$ for some $r \in I \cap h(R)$ and $s \in J \cap h(R)$. Since $rsK \subseteq N$ and $rs \notin (N :_R M)$, it follows from (vi) that $rK \subseteq N$ or $sK \subseteq N$. Let us consider three cases:

Case 1. Let $rK \subseteq N$ but $sK \not\subseteq N$. Since $asK \subseteq N$ but neither $aK \subseteq N$ nor $sK \subseteq N$, we conclude from (vi) that $as \in (N :_R M)$. Since $aK \not\subseteq N$ but $rK \subseteq N$, we conclude that $(a+r)K \not\subseteq N$. Now since $(a+r)sK \subseteq N$ but neither $(a+r)K \subseteq N$ nor $sK \subseteq N$, (vi) implies $(a+r)s = as + rs \in (N :_R M)$. Then $rs \in (N :_R M)$, which is a contradiction.

Case 2. Let $rK \not\subseteq N$ but $sK \subseteq N$. Hence the proof is the same as in **Case 1**.

Case 3. Let $rK \subseteq N$ and $sK \subseteq N$. Firstly, we consider that $rK \subseteq N$. Since $rK \subseteq N$ and $aK \not\subseteq N$, we have $(a+r)K \not\subseteq N$. Now since $(a+r)bK \subseteq N$ but neither $(a+r)K \subseteq N$ nor $bK \subseteq N$, it follows from (vi) that $(a+r)b = ab + rb \in (N :_R M)$. Then $rb \in (N :_R M)$. Now we consider that $sK \subseteq N$. Since $sK \subseteq N$ and $bK \not\subseteq N$, we have $(b+s)K \not\subseteq N$. Now since $a(b+s)K \subseteq N$ but neither $aK \subseteq N$ nor $(b+s)K \subseteq N$, from (vi) we have $a(b+s) = ab + as \in (N :_R M)$. Then $as \in (N :_R M)$. Now since $(a+r)(b+s)K \subseteq N$ but neither $(a+r)K \subseteq N$ nor $(b+s)K \subseteq N$, we can conclude that $(a+r)(b+s) = ab + as + rb + rs \in (N :_R M)$ and then $rs \in (N :_R M)$, which is a contradiction. Hence $IK \subseteq N$ or $JK \subseteq N$, as needed. \square

THEOREM 2.2. *Let R be a G -graded commutative ring, M be a graded R -module*

and N be a graded submodule of M . If N is a graded 2-absorbing submodule of M , then $(N :_R M)$ is a graded 2-absorbing ideal. The converse is true if M is a Gr -multiplication R -module.

Proof. Assume that N is a graded 2-absorbing submodule. Let $abc \in (N :_R M)$ for some $a, b, c \in h(R)$. Then $abcM \subseteq N$; and put $cM = K$ such that K is a graded submodule of M . Hence $abK \subseteq N$ and so, by Theorem 2.1 (vi), either $ab \in (N :_R M)$ or $aK \subseteq N$ (so $acM \subseteq N$) or $bK \subseteq N$ (so $bcM \subseteq N$). Then $ab \in (N :_R M)$ or $bc \in (N :_R M)$ or $ac \in (N :_R M)$, as needed.

Conversely, suppose that $abm \in N$ for some $a, b \in h(R)$ and $m \in h(M)$ with $ab \notin (N :_R M)$. Since M is a Gr -multiplication R -module, there exists a graded ideal I of $G(R)$ such that $m = IM$. Then $abIM \subseteq N$ and so $abI \subseteq (N :_R M)$. Since $(N :_R M)$ is a graded 2-absorbing ideal and $ab \notin (N :_R M)$, we claim that $aI \subseteq (N :_R M)$ or $bI \subseteq (N :_R M)$. Otherwise, neither $aI \subseteq (N :_R M)$ nor $bI \subseteq (N :_R M)$. Then there exists $i_1, i_2 \in I \cap h(R)$ such that $ai_1 \notin (N :_R M)$ and $bi_2 \notin (N :_R M)$. Since $abi_1 \in (N :_R M)$ but $ab \notin (N :_R M)$, $ai_1 \notin (N :_R M)$ and $(N :_R M)$ is a graded 2-absorbing ideal, we have $bi_1 \in (N :_R M)$. Similarly for the next term, since $abi_2 \in (N :_R M)$ but $ab \notin (N :_R M)$, $bi_2 \notin (N :_R M)$ and $(N :_R M)$ is a graded 2-absorbing ideal, we have $ai_2 \in (N :_R M)$. Now since $ab(i_1 + i_2) \in (N :_R M)$, $ab \notin (N :_R M)$ and $(N :_R M)$ is a graded 2-absorbing ideal, we conclude that $a(i_1 + i_2) \in (N :_R M)$ or $b(i_1 + i_2) \in (N :_R M)$. If $a(i_1 + i_2) = ai_1 + ai_2 \in (N :_R M)$, then $ai_1 \in (N :_R M)$, which is a contradiction. If $b(i_1 + i_2) = bi_1 + bi_2 \in (N :_R M)$, then $bi_2 \in (N :_R M)$, which is a contradiction. Thus $aI \subseteq (N :_R M)$ (so $aIM \subseteq N$) or $bI \subseteq (N :_R M)$ (so $bIM \subseteq N$). Hence $ab \in (N :_R M)$ or $am \in N$ or $bm \in N$. Therefore $(N :_R M)$ is a graded 2-absorbing ideal of R . \square

PROPOSITION 2.3. *Let R be a G -graded commutative ring, M be a graded R -module and N be a graded proper submodule of M . If $(N :_R M)$ is a graded 2-absorbing ideal, then $Gr(N :_R M)$ is a graded 2-absorbing ideal.*

Proof. Assume that $abc \in Gr(N :_R M)$ for some $a, b, c \in h(R)$ with $ab \notin Gr(N :_R M)$ and $bc \notin Gr(N :_R M)$. Then there exists a positive integer n such that $(abc)^n = a^n b^n c^n \in Gr(N :_R M)$ with $a^n b^n \notin (N :_R M)$ and $b^n c^n \notin (N :_R M)$. Since $(N :_R M)$ is a graded 2-absorbing ideal, we have $a^n c^n \in (N :_R M)$. Then $ac \in Gr(N :_R M)$. \square

The *graded radical* of a graded submodule N of a graded R -module M , which is denoted by $Gr_M(N)$, is defined to be the intersection of all graded prime submodules of M containing N . If N is not contained in any graded prime submodule of M , then $Gr_M(N) = M$.

THEOREM 2.4. *Let R be a G -graded commutative ring, M be a Gr -multiplication R -module and N be a graded submodule of M . If N is a graded 2-absorbing submodule of M , then $Gr_M(N)$ is a graded 2-absorbing submodule of M .*

Proof. Since N is a graded 2-absorbing submodule, by Theorem 2.2, $(N :_R M)$ is a graded 2-absorbing ideal of $G(R)$. Then $Gr(N :_R M)$ is a graded 2-absorbing ideal of $G(R)$, by Proposition 2.3. By [14, Lemma 2], $Gr(N :_R M) = (Gr_M(N) :_R M)$.

Hence $(Gr_M(N) :_R M)$ is a graded 2-absorbing ideal of $G(R)$. By Theorem 2.2, since M is a Gr -multiplication R -module, we get that $Gr_M(N)$ is a graded 2-absorbing submodule of M . \square

PROPOSITION 2.5. *Let R be a G -graded commutative ring and M be a Gr -multiplication R -module. Suppose that N and K are distinct graded prime submodules of M . Then the following statements hold:*

- (i) $(N :_R M) \cap (K :_R M)$ is a graded 2-absorbing ideal of $G(R)$;
- (ii) $N \cap K$ is a graded 2-absorbing submodule of M .

Proof. (i) Since N and K are graded prime submodules of M , by [4, Proposition 2.5], $(N :_R M)$ and $(K :_R M)$ are graded prime ideals of $G(R)$. Then $(N :_R M) \cap (K :_R M)$ is a graded 2-absorbing ideal of $G(R)$, by [13, Theorem 2.5].

(ii) Since N and K are graded prime submodules of M , by [4, Proposition 2.5], $(N :_R M)$ and $(K :_R M)$ are graded prime ideals of $G(R)$. Then (i) implies that $(N :_R M) \cap (K :_R M) = (N \cap K :_R M)$ is a graded 2-absorbing ideal, and hence $N \cap K$ is a graded 2-absorbing submodule by Theorem 2.2. \square

THEOREM 2.6. *Let R be a G -graded commutative ring and M be a graded R -module. Suppose that N_1, \dots, N_k are graded submodules of M . If N_1, \dots, N_k are graded 2-absorbing submodules of M , then $\bigcap_{i=1}^k N_i$ is a graded 2-absorbing submodule of M .*

Proof. Let $a, b \in h(R)$ and $m \in h(M)$ such that $abm \in \bigcap_{i=1}^k N_i$ with $ab \notin (\bigcap_{i=1}^k N_i :_R M) = \bigcap_{i=1}^k (N_i :_R M)$. Then there exists i such that $ab \notin (N_i :_R M)$. Since $abm \in N_i$, $ab \notin (N_i :_R M)$ and N_i is a graded 2-absorbing submodule of M , we conclude that $am \in N_i$ or $bm \in N_i$. Then $am \in \bigcap_{i=1}^k N_i$ or $bm \in \bigcap_{i=1}^k N_i$. Therefore $\bigcap_{i=1}^k N_i$ is graded 2-absorbing. \square

THEOREM 2.7. *Let R be a G -graded commutative ring, M be a graded R -module and N be a graded submodule of M . Suppose that N is a graded 2-absorbing submodule and $Gr(N :_R M) = P$ such that P is a graded prime ideal of $G(R)$. Then the following statements hold:*

- (i) If $m \in h(M) \setminus N$, then $Gr(N :_R m)$ is a graded prime ideal of $G(R)$ containing P ;
- (ii) If $m, m' \in h(M) \setminus N$, then either $Gr(N :_R m) \subseteq Gr(N :_R m')$ or $Gr(N :_R m') \subseteq Gr(N :_R m)$.

Proof. (i) Assume that $a, b \in h(R)$ such that $ab \in Gr(N :_R m)$. Then there exists some positive integer n such that $a^n b^n m \in N$. Since N is a graded 2-absorbing submodule of M , we conclude that $a^n b^n \in (N :_R M)$ or $a^n m \in N$ or $b^n m \in N$. If both these conditions hold, then we are done. Suppose that $a^n b^n \in (N :_R M)$. Since $a = \sum_{g \in G} a_g$ and $b = \sum_{g \in G} b_g$, and it is $a^n b^n = \sum_{g \in G} a_g^n b_g^n \in (N :_R M)$ for some smallest positive integer $n > 0$. Thus $ab \in Gr(N :_R M) = P$. Since P is a graded prime ideal, we have $a \in P$ or $b \in P$. Since $P = Gr(N :_R M) \subseteq Gr(N :_R m)$, we conclude that $a \in Gr(N :_R m)$ or $b \in Gr(N :_R m)$, as needed.

(ii) Assume that $Gr(N :_R m) \not\subseteq Gr(N :_R m')$. Let $a \in Gr(N :_R m)$ and $b \in Gr(N :_R m') \setminus Gr(N :_R m)$. Then there exists a smallest positive integer $n > 0$ such that $a_g^n m \in N$, $b_g^n m' \in N$ and $b_g^n m \notin N$ for $g \in G$. If $a_g^n(m + m') \in N$, then $a_g^n m' \in N$

for $g \in G$ and so $a^n m' = \sum_{g \in G} a_g^n m' \in N$. Thus $a \in Gr(N :_R m')$. Suppose that $a_g^n(m + m') \notin N$. Since N is a graded 2-absorbing submodule, $a_g^n b_g^n(m + m') \in N$ but $a_g^n(m + m') \notin N$ and $b_g^n(m + m') \notin N$, then $a^n b^n = \sum_{g \in G} a_g^n b_g^n \in (N :_R M)$ and so $ab \in Gr(N :_R M)$. Then $ab \in P$ where P is a graded prime ideal. Thus $a \in P$ or $b \in P$. If $b \in P$, then $b^n m \in N$, which is a contradiction. If $a \in P$, then $a \in Gr(N :_R m')$, as needed. \square

THEOREM 2.8. *Let R be a G -graded commutative ring, M be a graded R -module and N be a graded submodule of M . Suppose that N is a graded 2-absorbing submodule and $Gr(N :_R M) = P \cap Q$ such that P and Q are graded prime ideals of $G(R)$. Then the following statements hold:*

- (i) *If $m \in h(M) \setminus N$ and $P \subseteq Gr(N :_R m)$, then $Gr(N :_R m)$ is a graded prime ideal of $G(R)$;*
- (ii) *If $m, m' \in h(M) \setminus N$ and $P \subseteq Gr(N :_R m) \cap Gr(N :_R m')$, then either $Gr(N :_R m) \subseteq Gr(N :_R m')$ or $Gr(N :_R m') \subseteq Gr(N :_R m)$.*

Proof. The proof is similar to the proof of Theorem 2.7. \square

Let R be a G -graded ring, M be a graded R -module and N be a graded submodule of M . Recall that $(N :_M I) = \{m \in h(M) \mid mI \subseteq N\}$ and $(N :_M I^\infty)$ are graded submodules of graded R -module M .

LEMMA 2.9. *Let R be a G -graded commutative ring, M be a graded R -module and N be a graded submodule of M . Then $((N :_M I) :_R M) = ((N :_R M) :_M I)$.*

THEOREM 2.10. *Let R be a G -graded commutative ring, M be a graded R -module and N be a graded submodule of M . If N is a graded 2-absorbing submodule of M , then $(N :_M I)$ is a graded 2-absorbing submodule of M .*

Proof. Assume that $abm \in (N :_M I)$ for some $a, b \in h(R)$ and $m \in h(M)$. Then $Iabm \in N$ and hence, by Theorem 2.1 (iv), $Im \subseteq N$ or $abm \in N$ or $Iab \subseteq (N :_R N)$. If $Im \subseteq N$, then we are done by definition. If $Iab \subseteq (N :_R N)$, then by Lemma 2.9, $ab \in ((N :_R M) :_M I) = ((N :_M I) :_R M)$. If $abm \in N$, then $ab \in (N :_R M)$ or $am \in N$ or $bm \in N$. Since N is a graded 2-absorbing submodule, hence $Iab \subseteq (N :_R M)$ (so $ab \in ((N :_M I) :_R M)$) or $Iam \in N$ (so $am \in (N :_M I)$) or $Ibm \in N$ (so $bm \in (N :_M I)$). Therefore $(N :_M I)$ is a graded 2-absorbing submodule. \square

COROLLARY 2.11. *Let N be a graded submodule of graded R -module M . If N is a graded 2-absorbing submodule, then N_r is a graded 2-absorbing submodule for every $r \in h(R) \setminus (N :_M r)$. Moreover, $(N :_M I^n) = (N :_M I^{n+1})$ for all $n \geq 2$.*

Proof. Assume that $abm \in N_r = (N :_M r)$ for some $a, b \in h(R)$, $m \in h(M)$ and $r \in h(R) \setminus (N :_M r)$. Then $ab(rm) \in N$. Since N is a graded 2-absorbing submodule, we conclude that $ab \in (N :_R M)$ or $a(rm) \in N$ or $b(rm) \in N$. If both these conditions hold, then we are done. If $ab \in (N :_R M) \subseteq (N_r :_R M)$, then $ab \in (N_r :_R M)$. Therefore N_r is a graded 2-absorbing submodule of M for every $r \in h(R) \setminus (N :_M r)$. Now suppose that $m \in (N :_M I^3)$ for $m \in h(M)$. Then $I^3 m \subseteq N$. By Theorem 2.1 (v), $I^2 m \subseteq N$ or $Im \subseteq N$ or $I^3 \subseteq (N :_R M)$. If the first two

conditions hold, then we are done. If $I^3 \subseteq (N :_R M)$, then $I^2 \subseteq (N :_R M)$. Since $(N :_R M)$ is a graded 2-absorbing ideal, by Theorem 2.2, then $(N :_M I^3) \subseteq (N :_M I^2)$. The converse inclusion is obvious, so $(N :_M I^n) = (N :_M I^{n+1})$ for all $n \geq 2$. \square

THEOREM 2.12. *Let R be a G -graded commutative ring, M be a graded R -module and N be a graded submodule of M . Suppose that $Gr(N :_R M)$ is a graded prime ideal with $Gr(N :_R M) \neq (N :_R M)$. If $(N_r :_R M)$ is a graded prime ideal such that $r \in Gr(N :_R M) \setminus (N :_R M)$, then $(N :_R M)$ is a graded 2-absorbing ideal of $G(R)$.*

Proof. Assume that $rst \in (N :_R M)$ for some $r, s, t \in h(R)$ with $st \notin (N :_R M)$. Since $Gr(N :_R M)$ is a graded prime ideal, we may assume that $r \in Gr(N :_R M)$. If $r \in (N :_R M)$, then we are done. So we can suppose that $r \notin (N :_R M)$, and then $r \in Gr(N :_R M) \setminus (N :_R M)$. Since $(N_r :_R M)$ is a graded prime ideal and $rst \in (N_r :_R M)$, we conclude that $s \in (N_r :_R M)$ or $t \in (N_r :_R M)$. Hence $rs \in (N :_R M)$ or $rt \in (N :_R M)$. \square

THEOREM 2.13. *Let R be a G -graded commutative ring, M be a graded R -module and N be a graded submodule of M . Suppose that $Gr(N :_R m)$ is a graded prime ideal with $Gr(N :_R m) \neq (N :_R m)$ for all $m \in h(M) \setminus N$. If N_r is a graded prime submodule for $r \in Gr(N :_R m) \setminus (N :_R m)$, then N is a graded 2-absorbing submodule of M .*

Proof. Assume that $r, s \in h(R)$ and $m \in h(M)$ such that $rs m \in N$. Then $rs \in (N :_R m)$. Since $Gr(N :_R m)$ is a graded prime ideal, we may suppose that $r \in Gr(N :_R m)$. If $r \in (N :_R m)$, then we are done. Let $r \notin (N :_R m)$. Then $r \in Gr(N :_R m) \setminus (N :_R m)$. Since N_r is a graded prime submodule and $sm \in (N :_M r) = N_r$, we have $s \in (N_r :_R M)$ or $m \in N_r$ and so $rs \in (N :_R M)$ or $rm \in N$. Hence N is a graded 2-absorbing submodule of M . \square

THEOREM 2.14. *Let R be a G -graded commutative ring, M be a graded R -module and N be a submodule of M . Then the following statements are equivalent:*

- (i) N is a graded 2-absorbing submodule;
- (ii) For every graded ideal I, J of $G(R)$ and every graded submodule K of M with $(K + IL) \cap S \neq \emptyset$, $(K + JL) \cap S \neq \emptyset$ and $(K + IJM) \cap S \neq \emptyset$ such that $S = M \setminus N$, implies that $(K + IJL) \cap S \neq \emptyset$.

Proof. (i) \Rightarrow (ii) Let N be a graded 2-absorbing submodule. Suppose that I, J are graded ideals of $G(R)$ and K, L are graded submodules of M such that $(K + IL) \cap S \neq \emptyset$, $(K + JL) \cap S \neq \emptyset$ and $(K + IJM) \cap S \neq \emptyset$. If $(K + IJL) \cap S = \emptyset$, then $(K + IJL) \subseteq N$ and so $IJL \subseteq N$. Since N is a graded 2-absorbing submodule, we conclude that $IL \subseteq N$ or $JL \subseteq N$ or $IJM \subseteq N$, by Theorem 2.1 (ix). Then $(K + IL) \cap S = \emptyset$ or $(K + JL) \cap S = \emptyset$ or $(K + IJM) \cap S = \emptyset$, which are contradictions. Hence $(K + IJL) \cap S \neq \emptyset$, as needed.

(ii) \Rightarrow (i) Suppose that $IJL \subseteq N$ for some graded ideals I, J of $G(R)$ and some graded submodule L of M . We may assume that neither $IL \subseteq N$ nor $JL \subseteq N$ nor $IJM \subseteq N$. Then $(K + IL) \cap S \neq \emptyset$, $(K + JL) \cap S \neq \emptyset$ and $(K + IJM) \cap S \neq \emptyset$ and hence $(K + IJL) \cap S \neq \emptyset$, which is a contradiction. Therefore N is a graded 2-absorbing submodule of M . \square

3. Graded classical 2-absorbing submodules

In this section we will define the concept of graded classical 2-absorbing submodules as a generalization of graded classical prime submodules. Also, we show a number of results of graded classical 2-absorbing submodules.

DEFINITION 3.1. Let R be a G -graded ring and M be a graded R -module. A proper graded submodule N of M is called a *graded classical 2-absorbing submodule*, if whenever $a, b, c \in h(R)$ and $m \in h(M)$ with $abcm \in N$, then either $abm \in N$ or $bcm \in N$ or $acm \in N$. It is denoted as ${}^{gr}Cl(2)$ -absorbing submodule.

LEMMA 3.2. Let R be a G -graded ring and M be a graded R -module. Then the following statements hold:

- (i) Every graded classical prime submodule is a graded 2-absorbing submodule and a ${}^{gr}Cl(2)$ -absorbing submodule;
- (ii) If N is a graded classical prime submodule, then $(N :_R M)$ is a graded prime ideal, [2, Lemma 3.1].

In what follows, we show the basic theorem on ${}^{gr}Cl(2)$ -absorbing submodules.

THEOREM 3.3. Let R be a G -graded ring, M be a graded R -module and N be a proper graded submodule of M . Suppose that N is a ${}^{gr}Cl(2)$ -absorbing submodule. Then the following statements hold:

- (i) For all elements $a, b, c \in h(R)$ either $(N :_M abc) = (N :_M ab)$ or $(N :_M abc) = (N :_M bc)$ or $(N :_M abc) = (N :_M ac)$;
- (ii) For all elements $a, b, c \in h(R)$ and every graded submodule K of M , $abcK \subseteq N$ implies that $abK \subseteq N$ or $bcK \subseteq N$ or $acK \subseteq N$;
- (iii) For all elements $a, b \in h(R)$ and every graded submodule K of M , $abK \not\subseteq N$ implies that $(N :_R abK) = (N :_R aK)$ or $(N :_R abK) = (N :_R bK)$;
- (iv) For all elements $a, b \in h(R)$, every graded ideal I of $G(R)$ and every graded submodule K of M , $abIK \subseteq N$ implies that $abK \subseteq N$ or $aIK \subseteq N$ or $bIK \subseteq N$;
- (v) For every element $a \in h(R)$, every graded ideal I of $G(R)$ and every graded submodule K of M , $aIK \not\subseteq N$ implies that $(N :_R aIK) = (N :_R aK)$ or $(N :_R aIK) = (N :_R IK)$;
- (vi) For every element $a \in h(R)$, every graded ideal I, J of $G(R)$ and every graded submodule K of M , $aIJK \subseteq N$ implies that $aIK \subseteq N$ or $aJK \subseteq N$ or $IJK \subseteq N$;
- (vii) For every graded ideal I, J of $G(R)$ and every graded submodule K of M , $IJK \not\subseteq N$ implies that $(N :_R IJK) = (N :_R IK)$ or $(N :_R IJK) = (N :_R JK)$;
- (viii) For every graded ideal I, J, P of $G(R)$ and every graded submodule K of M , $IJPK \subseteq N$ implies that $IJK \subseteq N$ or $JPK \subseteq N$ or $IPK \subseteq N$.

Proof. (i) Assume that $abcm \in N$ for some $m \in h(M)$. Since N is a ${}^{gr}Cl(2)$ -absorbing submodule, we have $abm \in N$ or $bcm \in N$ or $acm \in N$. Then either $m \in (N :_M ab)$ or $m \in (N :_M bc)$ or $m \in (N :_M ac)$ and hence $(N :_M abc) = (N :_M ab)$ or $(N :_M abc) = (N :_M bc)$ or $(N :_M abc) = (N :_M ac)$.

(ii) Assume that $abcK \subseteq N$ for some $a, b, c \in h(R)$ and some graded submodule K of M . Then $K \subseteq (N :_M abc)$ and so either $K \subseteq (N :_M ab)$ or $K \subseteq (N :_M bc)$ or

$K \subseteq (N :_M ac)$, it follows from (i). Hence either $abK \subseteq N$ or $bcK \subseteq N$ or $acK \subseteq N$, as needed.

(iii) Assume that K is a graded submodule of M and $a, b \in h(R)$ are such that $abK \not\subseteq N$. Let $x \in (N :_R abK)$ for some $x \in h(R)$ and thus $xabK \subseteq N$. Since $abK \not\subseteq N$, we have $xaK \subseteq N$ or $xbK \subseteq N$, what follows from (ii). Then either $x \in (N :_R aK)$ or $x \in (N :_R bK)$ and hence $(N :_R abK) = (N :_R aK)$ or $(N :_R abK) = (N :_R bK)$.

(iv) Assume that $abIK \subseteq N$ for some $a, b \in h(R)$, some graded ideal I and some graded submodule K of M ; thus $I \subseteq (N :_R abK)$. If $abK \subseteq N$, then we are done, what follows from (iii). If $abK \not\subseteq N$, then $I \subseteq (N :_R aK)$ or $I \subseteq (N :_R bK)$, what follows from (iii). Hence either $aIK \subseteq N$ or $bIK \subseteq N$, as required.

The proof of the remaining parts is similar to the previous ones, so we omit it. \square

THEOREM 3.4. *Let R be a G -graded ring, M be a graded R -module and N be a proper graded submodule of M . Then N is a ${}^{gr}Cl(2)$ -absorbing submodule of M if and only if for every graded submodule K of M such that $K \not\subseteq N$, $(N :_R K)$ is a graded 2-absorbing ideal.*

Proof. Assume that N is a ${}^{gr}Cl(2)$ -absorbing submodule of M . Let $a, b, c \in h(R)$ be such that $abc \in (N :_R K)$. Then $abcK \subseteq N$ and so $abK \subseteq N$ or $bcK \subseteq N$ or $acK \subseteq N$, by Theorem 3.3 (ii). Hence either $ab \in (N :_R K)$ or $bc \in (N :_R K)$ or $ac \in (N :_R K)$. Therefore $(N :_R K)$ is a graded 2-absorbing ideal of $G(R)$. Conversely, suppose that $abcL \subseteq N$ for some $a, b, c \in h(R)$ and some graded submodule L of M . If $L \subseteq N$, then we are done. If $L \not\subseteq N$, then $abc \in (N :_R L)$. Since $(N :_R L)$ is a graded 2-absorbing ideal, we conclude that $ab \in (N :_R L)$ or $bc \in (N :_R L)$ or $ac \in (N :_R L)$. Then either $abm \in N$ or $bcm \in N$ or $acm \in N$ for some $m \in h(M) \cap L$, as needed. \square

COROLLARY 3.5. *Let R be a G -graded ring, M be a graded R -module and N be a proper graded submodule of M . If N is a ${}^{gr}Cl(2)$ -absorbing submodule of M , then $(N :_R M)$ is a graded 2-absorbing ideal of $G(R)$. Moreover, for every $m \in h(M) \setminus N$, $(N :_R m)$ is a graded 2-absorbing ideal of $G(R)$.*

THEOREM 3.6. *Let R be a G -graded ring, M be a graded R -module and N be a graded proper submodule of M . If N is a graded 2-absorbing submodule of M , then N is a ${}^{gr}Cl(2)$ -absorbing submodule. The converse is true if M is a Gr -multiplication R -module.*

Proof. Assume that N is a graded 2-absorbing submodule. Let $a, b, c \in h(R)$ and $m \in h(M)$ be such that $abcm \in N$. Since N is a graded 2-absorbing submodule, we conclude that $ab \in (N :_R M)$ or $bcm \in N$ or $acm \in N$. If both cases are true, then we are done. If the first case holds, then $abm \in N$ and so N is ${}^{gr}Cl(2)$ -absorbing. Conversely, suppose that N is a ${}^{gr}Cl(2)$ -absorbing submodule of M . Then $(N :_R M)$ is a graded 2-absorbing ideal of $G(R)$, by Corollary 3.5. Since M is a Gr -multiplication R -module, by Theorem 2.2, N is a graded 2-absorbing submodule. \square

THEOREM 3.7. *Let R be a G -graded ring, M be a graded R -module and N be a proper graded submodule of M . Suppose that N is a ${}^{gr}Cl(2)$ -absorbing submodule. Then the following statements hold:*

- (i) For all elements $a, b, c \in h(R)$ either $(N :_M abc) = (N :_M ab)$ or $(N :_M abc) = (N :_M bc)$ or $(N :_M abc) = (N :_M ac)$;
- (ii) For all elements $a, b \in h(R)$ and $m \in h(M)$, $abm \not\subseteq N$ implies that $(N :_R abm) = (N :_R am)$ or $(N :_R abm) = (N :_R bm)$;
- (iii) For all elements $a, b \in h(R)$, $m \in h(M)$ and every graded ideal I of $G(R)$, $abIm \subseteq N$ implies that $abm \subseteq N$ or $aIm \subseteq N$ or $bIm \subseteq N$;
- (iv) For all elements $a \in h(R)$, $m \in h(M)$ and every graded ideal I of $G(R)$, $aIm \not\subseteq N$ implies that $(N :_R aIm) = (N :_R am)$ or $(N :_R aIm) = (N :_R Im)$;
- (v) For all elements $a \in h(R)$, $m \in h(M)$ and all graded ideals I, J of $G(R)$, $aIJm \subseteq N$ implies that $aIm \subseteq N$ or $aJm \subseteq N$ or $IJm \subseteq N$;
- (vi) For all graded ideals I, J of $G(R)$ and every element of submodule $m \in h(M)$, $IJm \not\subseteq N$ implies that $(N :_R IJm) = (N :_R Im)$ or $(N :_R IJm) = (N :_R Jm)$;
- (vii) For all graded ideals I, J, P of $G(R)$ and every element $m \in h(M)$, $IJPm \subseteq N$ implies that $IJm \subseteq N$ or $JPm \subseteq N$ or $IPm \subseteq N$.

Proof. The complete proof is similar to the proof of Theorem 3.3. \square

THEOREM 3.8. *Let R be a G -graded ring, M be a Gr -multiplication R -module and N be a graded submodule of M . Then the following statements are equivalent:*

- (i) N is a ${}^{gr}Cl(2)$ -absorbing submodule;
- (ii) For all graded submodules N_1, N_2 and N_3 of M and every element $m \in h(M)$, $N_1N_2N_3m \in N$ implies that $N_1N_2m \subseteq N$ or $N_2N_3m \subseteq N$ or $N_1N_3m \subseteq N$.

Proof. (i) \Rightarrow (ii) Suppose that $N_1N_2N_3m \in N$ for some graded submodules N_1, N_2 and N_3 of M and some $m \in h(M)$. Since M is a Gr -multiplication R -module, there exist graded ideals I_1, I_2 and I_3 of $G(R)$ such that $N_1 = I_1M$, $N_2 = I_2M$ and $N_3 = I_3M$. Then $I_1I_2I_3m \in N$ and so, by Theorem 3.7 (vii), either $I_1I_2m \subseteq N$ or $I_2I_3m \subseteq N$ or $I_1I_3m \subseteq N$. Hence either $N_1N_2m \subseteq N$ or $N_2N_3m \subseteq N$ or $N_1N_3m \subseteq N$.

(ii) \Rightarrow (i) Assume that $I_1I_2I_3m \in N$ for some graded ideals I_1, I_2 and I_3 of $G(R)$ and some $m \in h(M)$. Since M is a Gr -multiplication R -module, we obtain that $I_1M = N_1$, $N_2 = I_2M$ and $N_3 = I_3M$. Then the conclusion follows from (ii). \square

THEOREM 3.9. *Let R be a G -graded ring, M be a graded R -module and N be a submodule of M . Then the following statements are equivalent:*

- (i) N is a ${}^{gr}Cl(2)$ -absorbing submodule;
- (ii) For all graded ideals I, J, P of $G(R)$ and all graded submodules K, L of M , with $(K + IJL) \cap S \neq \emptyset$, $(K + JPL) \cap S \neq \emptyset$ and $(K + IPL) \cap S \neq \emptyset$, $S = M \setminus N$, implies that $(K + IJPL) \cap S \neq \emptyset$.

Proof. (i) \Rightarrow (ii) Assume that N is a ${}^{gr}Cl(2)$ -absorbing submodule. Suppose that I, J, P are graded ideals and K, L are graded submodules such that $(K + IJL) \cap S \neq \emptyset$, $(K + JPL) \cap S \neq \emptyset$ and $(K + IPL) \cap S \neq \emptyset$. Let $(K + IJPL) \cap S = \emptyset$. Then $K + IJPL \subseteq N$ and so $K \subseteq N$ and $IJPL \subseteq N$. Since N is a ${}^{gr}Cl(2)$ -absorbing submodule, Theorem 3.3 (viii) implies that $IJL \subseteq N$ or $JPL \subseteq N$ or $IPL \subseteq N$. Suppose that $IJL \subseteq N$, then $(K + IJL) \cap S = \emptyset$, which is a contradiction. In the next two cases we can obtain a contradiction in a similar way.

(ii) \Rightarrow (i) Suppose that $IJPL \subseteq N$ for some graded ideals I, J, P of $G(R)$ and some graded submodules K, L of M . If neither $IJL \subseteq N$ nor $JPL \subseteq N$ nor $IPL \subseteq N$,

then $IJL \cap S \neq \emptyset$, $JPL \cap S \neq \emptyset$ and $IPL \cap S \neq \emptyset$ and thus $IJPL \cap S \neq \emptyset$, which is a contradiction. Hence either $IJL \subseteq N$ or $JPL \subseteq N$ or $IPL \subseteq N$. Therefore N is a ${}^{gr}Cl(2)$ -absorbing submodule. \square

ACKNOWLEDGEMENT. The authors would like to thank the referees for their valuable comments and suggestions on the earlier version of the manuscript which improved it a lot.

REFERENCES

- [1] K. Al-Zoubi, R. Abu-Dawwas, *On graded 2-absorbing and weakly graded 2-absorbing submodules*, J. Math. Sciences: Advances and Applications., **28** (2014), 45–60.
- [2] K. AL-Zoubi, M. Jaradat, R. Abu-Dawwas, *On graded classical prime and graded prime submodules*, Bull. Iranian Math. Soc., **41** (1) (2015), 217–225.
- [3] S. E. Atani, *On graded prime submodules*, Chiang Mai J. Sci., **33** (1) (2006), 3–7.
- [4] S. E. Atani, F. Farzalipour, *On graded secondary modules*, Turk. J. Math., **31** (2007), 371–378.
- [5] A. Badawi, *On 2-absorbing ideals of commutative rings*, Bull. Austral. Math. Soc., **75** (2007), 417–429.
- [6] M. Baziar, M. Behboodi, *Classical primary submodules and decomposition theory of modules*, J. Algebra Appl., **8** (3) (2009), 351–369.
- [7] M. Behboodi, *A generalization of Baer’s lower nilradical for modules*, J. Algebra Appl., **6** (2) (2007), 337–353.
- [8] K. Hakan Oral, Ü. Tekir, A. G. Ağargün, *On graded prime and primary submodules*, Turk. J. Math., **35** (2011), 159–167.
- [9] H. Mostafanasab, E. Yetkin, Ü. Tekir, K.S.H. Oral, *Classical 2-absorbing submodules of modules over commutative rings*, Eur. J. Pure Appl. Math., **8** (3) (2015), 417–430.
- [10] C. Nastasescu, F. Van Oystaeyen, *Graded Ring Theory*, Mathematical Library 28, North Holland, Amsterdam, 1982.
- [11] S. Park, *Some properties of Gr-multiplication modules*, Korean J. Math., **20** (3) (2012), 315–321.
- [12] M. Refai, M. Hailat, S. Obiedat, *Graded radicals and graded prime spectra*, Far East J. Math. Sci., part I (2000), 59–73.
- [13] F. Soheilnia, A. Yousefian Darani, *On graded 2-absorbing primary and graded weakly 2-absorbing primary ideals*, Kyugpook Math. J., **57** (2017), 559–580.
- [14] F. Soheilnia, A. Yousefian Darani, *On graded 2-absorbing primary submodules of multiplication modules*, Preprint.
- [15] A. Yousefian Darani, F. Soheilnia, *On 2-absorbing and weakly 2-absorbing submodules*, Thai J. Math., **9** (2011), 577–584.
- [16] A. Yousefian Darani, F. Soheilnia, *On n-absorbing submodules*, Math. Commun., **17** (2012), 547–557.

(received 25.12.2017; in revised form 09.06.2018; available online 28.07.2018)

Department of Mathematics, Imam Khomeini International University, Qazvin, Iran

E-mail: soheilnia@gmail.com

Department of Mathematics and Applications, University of Mohaghegh Ardabili, 5619911367, Ardabil, Iran

E-mail: yousefian@uma.ac.ir