

## EQUIPRIME FUZZY GRAPH OF A NEARRING WITH RESPECT TO A LEVEL IDEAL

**B. Jagadeesha, Babushri Srinivas Kedukodi and Syam Prasad Kuncham**

**Abstract.** In this paper, we introduce an equiprime fuzzy graph of a nearring with respect to the level ideal of a fuzzy ideal. We interrelate graph theoretical properties of the graph and ideal theoretical properties of nearring. We show that the properties like vertex cut, connectedness of the graph depend on the properties of the fuzzy ideal. We define ideal symmetry of the graph and find conditions for the graph to be ideal symmetric. If the fuzzy ideal is equiprime then we show that the level set induces a fuzzy clique. We find conditions required for the level set to be the vertex cover of the graph. We find interrelation between equiprime fuzzy graph and fuzzy graph of nearring with respect to level ideal. We study properties of the graph under nearring homomorphism. We prove that the connectedness of the graph in homomorphic image depends on properties of ideal. We obtain conditions required for homomorphic image of an equiprime fuzzy ideal to be an equiprime fuzzy ideal.

### 1. Introduction

Algebra and graph theory are independent areas of mathematics, and researchers have attempted to link these areas. The study of the relation between algebraic structures and graph theory was initiated by Beck [6] by introducing the zero divisor graph of a commutative ring. Redmond [25] generalized the zero divisor graph using an ideal and defined an ideal-based zero divisor graph of a commutative ring. Bhavanari, Kuncham and Kedukodi [8] proposed the graph of a generalized ring (nearring) with respect to an ideal and introduced a new type of symmetry of the graph called ideal symmetry. Pang, Zhang, Zhang and Wang [23] studied the problem of dominating set in directed graphs.

Fuzzy set theory is applied in various areas of mathematics such as algebra, graph theory, topology, and differential equations to generalize various constraints in these areas. Bhakat and Das [7] used fuzzy set theory to generalize the definition of the ideal and introduced the fuzzy ideal of a ring. Davvaz [10] generalized fuzzy ideals and

---

*2020 Mathematics Subject Classification:* 16Y30, 03E72, 16Y99

*Keywords and phrases:* Nearring; equiprime; ideal; fuzzy graph; homomorphism.

introduced threshold-based fuzzy ideals. Kedukodi, Kuncham, and Bhavanari [18] studied threshold-based equiprime, 3-prime, and c-prime fuzzy ideals of nearrings and used them to study roughness in rings. Akram [2] generalized the definition of fuzzy ideals using triangular norms by introducing T-fuzzy ideals of nearrings and studied Artinian and Noetherian nearrings. Shum and Akram [26] determined properties of intuitionistic (T,S)-fuzzy ideals of a nearring and found their properties. Lattice is an algebraic structure useful for studying order between different sets and between elements of a set. To study ordered fuzzy sets, Goguen [11] introduced lattice valued fuzzy (L-fuzzy sets). Interval valued fuzzy sets are another generalization of fuzzy set theory where the fuzzy membership function is represented by a closed subinterval of  $[0, 1]$ . Davvaz [9] used interval-valued fuzzy sets and L-fuzzy sets to begin the study of interval-valued L-fuzzy ideals of nearrings. Jagadeesha, Kedukodi, and Kuncham [15, 16, 19] studied various types of fuzzy ideals of nearrings using triangular norms and interval-valued L-fuzzy sets.

Researchers used fuzzy set theory to generalize graph theory. Mordeson and Peng [22] introduced basic operations such as Cartesian product, union, intersection on fuzzy subgraphs. Koczy [20] introduced the concept of vertex fuzzy graph and applied it to distributed communication switching systems. Akram and Nawaz [4] related soft set theory and fuzzy graph theory by introducing fuzzy soft graphs. Akram and Dudek [3] related interval-valued fuzzy set theory and graph theory by introducing interval-valued fuzzy graphs. Akram [1] introduced a new type of fuzzy graphs, namely m-polar fuzzy graphs, and applied them to image processing and decision making.

In this paper, we relate equiprime fuzzy ideal of a nearring with a fuzzy graph. We introduce equiprime fuzzy graph of nearring  $N$  with respect to the level ideal  $\nu_t$  of a fuzzy ideal  $\nu$  denoted by  $(N_e, \nu, \rho_e, t)$ . We study the properties of the fuzzy graph. If  $x$  is an element of the fuzzy ideal, then we prove that  $x$  is connected to all other vertices of the fuzzy graph. If the fuzzy ideal is an equiprime fuzzy ideal, then the level set  $\nu_t$  is a strong vertex cut of the fuzzy graph. We define the ideal symmetry of the graph and prove that if  $\nu$  is an equiprime fuzzy ideal, then the fuzzy graph is ideal symmetric. We find conditions for a fuzzy clique of the fuzzy graph. If  $\nu$  is an equiprime fuzzy ideal, then we prove that  $\nu_t$  is the vertex cover of the fuzzy graph. If nearring is a simple nearring and the fuzzy ideal is an equiprime fuzzy ideal, then we prove that the fuzzy graph is a complete subgraph or a star graph. We find conditions under which the fuzzy graphs of nearring, c-prime fuzzy graph and equiprime fuzzy graph defined by Kedukodi, Kuncham and Bhavanari [17] are equivalent. If a fuzzy ideal is c-prime, then we prove that the fuzzy graph and the c-prime fuzzy graph are identical. If  $N_1$  and  $N_2$  are nearrings and  $f$  is a nearring homomorphism between  $N_1, N_2$ , then  $f$  is a graph homomorphism between  $(N_{1e}, \nu, \rho_e, t)$  and  $(N_{2e}, f(\nu), f(\rho)_e, t)$ . We prove that if  $x$  is connected to all vertices of  $(N_{1e}, \nu, \rho_e, t)$ , then  $f(x)$  is connected to all other vertices of  $(N_{2e}, f(\nu), f(\rho)_e, t)$ . We find conditions necessary for a homomorphic image of an equiprime fuzzy ideal to be an equiprime fuzzy ideal.

## 2. Preliminaries

In this paper  $N, N_1$  and  $N_2$  represent right nearrings. We refer to Pilz [24] for definitions, concepts related to nearrings, Anderson and Fuller [5] for rings, Harary [13] for graphs.

DEFINITION 2.1 ([14]). Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs. A *graph homomorphism* from  $G_1$  to  $G_2$  is a mapping  $f : V_1 \rightarrow V_2$  such that  $(f(u), f(v)) \in E_2$  whenever  $(u, v) \in E_1$ .

DEFINITION 2.2 ([21]). A fuzzy graph  $H = (\nu, \rho)$  of  $(V, E)$  is defined by a fuzzy subset  $\nu$  of  $V$  and a fuzzy subset  $\rho$  of  $E$  such that  $\rho(x, y) \leq \nu(x) \wedge \nu(y) \forall x, y \in V$ .

DEFINITION 2.3 ([21]). Let  $t \in [0, 1]$ . Then  $\nu_t = \{x \in V \mid \nu(x) \geq t\}$  is called the level set of  $\nu$ . Also  $\rho_t = \{(x, y) \in E \mid \rho(x, y) \geq t\}$  is called the level set of  $\rho$ . Then  $(\nu_t, \rho_t)$  is a graph with vertex set  $\nu_t$  and edge set  $\rho_t$ . Let  $(\nu, \rho)$  be a fuzzy graph. Then  $(\nu, \rho)$  is said to be *complete* if  $\rho(u, v) = \nu(u) \wedge \nu(v) \forall u, v \in V$ . A complete fuzzy subgraph with maximum number of vertices is called a *fuzzy clique* of  $(\nu, \rho)$ .

DEFINITION 2.4 ([24]). Let  $I, J, K$  be ideals of  $N$ . Then  $I$  is said to be a *prime ideal* if  $JK \subseteq I$  implies  $J \subseteq I$  or  $K \subseteq I$ . Nearring  $N$  is said to be *integral* if it has no zero divisors. Nearring  $N$  is said to be a *zerosymmetric* if  $x0 = 0$  for all  $x \in N$ . An ideal  $I$  of  $N$  is said to be *totally reflexive* if  $aNb \subseteq I$  then  $bNa \subseteq I$  for all  $a, b \in N$ . Nearring  $N$  is said to be *simple* if its only ideals are  $\{0\}$  and  $N$ . Let  $N_1, N_2$  be nearrings. Then  $g : N_1 \rightarrow N_2$  is said to be a *nearring homomorphism* if for all  $a, b \in N_1$ , (i)  $g(a + b) = g(a) + g(b)$ , (ii)  $g(ab) = g(a)g(b)$ .

DEFINITION 2.5 ([12]). Let  $I$  be an ideal of  $N$ . Then  $I$  is said to be a *c-prime ideal* of  $N$  if  $xy \in I$  implies  $x \in I$  or  $y \in I$  for all  $x, y \in N$ .  $I$  is said to be an *equiprime ideal* if  $a, x, y \in N$  with  $anx - any \in I$  for all  $n \in N$  implies  $a \in I$  or  $x - y \in I$ .  $I$  is said to be a *3-prime ideal* if  $x, y \in N$  and  $xny \in I$  for all  $n \in N$  implies  $x \in I$  or  $y \in I$ .

THEOREM 2.6 ([12, 27]). *The following implications hold in nearrings:*

(i) *each equiprime ideal is a 3-prime ideal;*

(ii) *each c-prime ideal is a 3-prime ideal.*

*The notions of equiprime ideal, 3-prime ideal and prime ideal coincide in rings. In commutative rings, all the above notions coincide.*

DEFINITION 2.7 ([12]). An ideal  $I$  of  $N$  is said to be a *c-semiprime ideal* if  $x \in N$  and  $x^2 \in I$  implies  $x \in I$ . An ideal  $I$  of  $N$  is said to be an *equisemiprime ideal* if, for  $a, b \in N$ ,  $(a - b)ra - (a - b)rb \in I$  for all  $r \in N$  implies  $(a - b) \in I$ .

DEFINITION 2.8 ([10]). Let  $\alpha, \beta \in [0, 1]$  and  $\alpha < \beta$ . Let  $\nu$  be a fuzzy subset of  $N$ . Then  $\nu$  is called a fuzzy ideal with thresholds  $\alpha, \beta$  if for all  $x, y, i \in N$ ,

(i)  $\alpha \vee \nu(x + y) \geq \beta \wedge \nu(x) \wedge \nu(y)$ , (ii)  $\alpha \vee \nu(-x) \geq \beta \wedge \nu(x)$ ,

(iii)  $\alpha \vee \nu(y + x - y) \geq \beta \wedge \nu(x)$ , (iv)  $\alpha \vee \nu(xy) \geq \beta \wedge \nu(x)$ ,

(v)  $\alpha \vee \nu(x(y + i) - xy) \geq \beta \wedge \nu(i)$ .

Here  $\alpha$  is called the lower threshold of  $\nu$  and  $\beta$  as the upper threshold of  $\nu$ . In this paper fuzzy ideal  $\nu$  means fuzzy ideal with lower threshold  $\alpha$  and upper threshold  $\beta$ .

**THEOREM 2.9** ([10]). *Let  $\nu$  be a fuzzy subset of  $N$ . Then  $\nu$  is a fuzzy ideal of  $N$  if and only if for every  $t \in (\alpha, \beta]$  the level subset  $\nu_t$  is an ideal of  $N$ .*

**DEFINITION 2.10** ([18]). Let  $\alpha, \beta \in [0, 1]$  and  $\alpha < \beta$ . Let  $\nu$  be a fuzzy ideal with thresholds  $\alpha, \beta$ . If  $x, y, a, b \in N$  then

(i)  $\nu$  is called an *equiprime fuzzy ideal with thresholds  $\alpha, \beta$*  if  $\alpha \vee \nu(a) \vee \nu(x - y) \geq \beta \wedge \inf_{r \in N} \nu(arx - ary)$ .

(ii)  $\nu$  is called a *3-prime fuzzy ideal with thresholds  $\alpha, \beta$*  if  $\alpha \vee \nu(a) \vee \nu(b) \geq \beta \wedge \inf_{r \in N} \nu(arb)$ .

(iii)  $\mu$  is called a *c-prime fuzzy ideal with thresholds  $\alpha, \beta$*  if  $\alpha \vee \nu(a) \vee \nu(b) \geq \beta \wedge \nu(ab)$ .

(iv) A fuzzy ideal  $\mu$  is called an *equisemiprime fuzzy ideal* if for all  $a \in N$ ,  $\alpha \vee \nu(a - b) \geq \beta \wedge \inf_{r \in N} \nu((a - b)ra - (a - b)rb)$ .

**THEOREM 2.11** ([18]). *Let  $\nu$  be a fuzzy ideal of  $N$ . Then  $\nu$  is an equiprime (resp. equisemiprime) fuzzy ideal of  $N$  if and only if for every  $t \in (\alpha, \beta]$ , the level subset  $\nu_t$  is a equiprime (resp. equisemiprime) ideal of  $N$ .*

**DEFINITION 2.12** ([9]). Let  $f : N_1 \rightarrow N_2$  be a mapping. Let  $\mu$  be a fuzzy subset of  $N_1$ . Then the image of  $\mu$  under  $f$  is given by

$$f(\hat{\mu})(y) = \begin{cases} \bigvee_{x \in f^{-1}(y)} \hat{\mu}(x) & \text{if } f^{-1}(y) \neq \emptyset \\ 0 & \text{if } f^{-1}(y) = \emptyset \end{cases}$$

for all  $y \in N_2$ .

**DEFINITION 2.13** ([17]). Let  $\nu : N \rightarrow (0, 1]$  be a fuzzy ideal of  $N$ . Let  $t \in (\alpha, \beta]$  be fixed. Define  $\rho : N \times N \rightarrow [0, 1]$  as follows:

$$\rho_c(x, y) = \begin{cases} \nu(x) \wedge \nu(y) & x \neq y \text{ and } (xNy \subseteq \nu_t \text{ or } yNx \subseteq \nu_t) \\ 0 & \text{otherwise.} \end{cases}$$

Then the fuzzy graph with respect to  $\nu_t$  denoted by  $(N_3, \nu, \rho_3, t)$  is called *fuzzy graph of  $N$  with respect to level ideal  $\nu_t$* .

**DEFINITION 2.14.** Let  $\nu : N \rightarrow (0, 1]$  be a fuzzy ideal of  $N$ . Let  $t \in (\alpha, \beta]$  be fixed. Define  $\rho : N \times N \rightarrow [0, 1]$  as follows:

$$\rho_c(x, y) = \begin{cases} \nu(x) \wedge \nu(y) & x \neq y \text{ and } (xy \in \nu_t \text{ or } yx \in \nu_t) \\ 0 & \text{otherwise} \end{cases}$$

Then the fuzzy graph with respect to  $\nu_t$  denoted by  $(N_c, \nu, \rho_c, t)$  is called *c-prime fuzzy graph of  $N$  with respect to level ideal  $\nu_t$* .

**REMARK 2.15.** In the next part of this paper by fuzzy (resp. fuzzy equiprime, fuzzy equisemiprime) ideal we mean fuzzy (resp. fuzzy equiprime, fuzzy equisemiprime) ideal with lower threshold  $\alpha$  upper threshold  $\beta$ .

### 3. Equiprime fuzzy graph of nearring with respect to level ideal

DEFINITION 3.1. Let  $\nu : N \rightarrow (0, 1]$  be a fuzzy ideal of  $N$ . Let  $t \in (\alpha, \beta]$  be fixed. For  $p \in N$ , define  $\rho_p : N \times N \rightarrow [0, 1]$  as follows:

$$\rho_p(p, x) = \begin{cases} \nu(p) \wedge \nu(x) & \text{if } p \neq x \text{ and } (prx - pr0 \in \nu_t) \\ & \text{or } (xrp - xr0 \in \nu_t) \text{ for all } r \in N \\ 0 & \text{otherwise.} \end{cases}$$

Then the fuzzy graph  $(N, \nu, \rho_p)$  is called the fuzzy graph of  $N$  with respect to  $p$  and level ideal  $\nu_t$ . We denote this fuzzy graph by  $(N_e, \nu, \rho_p, t)$ .

Let  $(N, \nu, \rho) = \bigcup_{p \in N} (N, \nu, \rho_p)$ . Then the fuzzy graph  $(N, \nu, \rho)$  is called the *equiprime fuzzy graph of  $N$  with respect level ideal  $\nu_t$* . We denote this fuzzy graph by  $(N_e, \nu, \rho_e, t)$

Now we provide examples for equiprime fuzzy graph of nearring with respect to level ideal.

EXAMPLE 3.2. Let  $N = \{0, a, b, c\}$  be the nearring with addition and multiplication defined as in Table 1.

+	0	a	b	c
0	0	a	b	c
a	a	0	c	b
b	b	c	0	a
c	c	b	a	0

·	0	a	b	c
0	0	0	0	0
a	0	a	0	a
b	b	b	b	b
c	b	c	b	c

Table 1: Nearing for Example 3.2

We define  $\nu : N \rightarrow (0, 1]$  by



$$\nu(x) = \begin{cases} 0.8 & \text{if } x = 0 \\ 0.5 & \text{if } x = b \\ 0.1 & \text{if } x \in \{a, c\}. \end{cases}$$

If we take thresholds  $\alpha = 0.1$  and  $\beta = 0.5$  then  $\nu$  is a fuzzy ideal of  $N$ . Let  $t = \beta$ . Then  $\nu_t = \{0, b\}$  and the values of  $\rho_p(p, x)$  are given in Table 2.

$\rho_p(p, x)$	$x = 0$	$x = a$	$x = b$	$x = c$
$\rho_0(0, x)$	0	0.1	0.5	0.1
$\rho_a(a, x)$	0.1	0	0.1	0
$\rho_b(b, x)$	0.5	0.1	0	0.1
$\rho_c(c, x)$	0.1	0	0.1	0

Table 2:  $\rho_p(p, x)$  when  $\nu_t = \{0, b\}$

Graphs  $(N_e, \nu, \rho_0, t)$ ,  $(N_e, \nu, \rho_a, t)$ ,  $(N_e, \nu, \rho_b, t)$ ,  $(N_e, \nu, \rho_c, t)$  are shown in Figures 1a, 1b, 1c and 1d respectively and the graph  $(N_e, \nu, \rho_e, t)$  is shown on Figure 2.

Figure 1:  $EQ_t^p(N)$ Figure 2:  $(N_e, \nu, \rho_e, t)$ 

EXAMPLE 3.3. Let  $N = Z_8 = \{0, 1, 2, 3, 4, 5, 6, 7\}$  be the ring of integers modulo 8. We define

$$\nu(x) = \begin{cases} 0.9 & \text{if } x = 0, \\ 0.7 & \text{if } x = 4, \\ 0.3 & \text{if } x \in \{2, 6\}, \\ 0.1 & \text{otherwise.} \end{cases}$$

If we take thresholds  $\alpha = 0.1$  and  $\beta = 0.3$  then  $\nu$  is a fuzzy ideal of  $N$ . Let  $t = \beta$ . Then  $\nu_t = \{0, 2, 4, 6\}$  and the values of  $\rho_p(p, x)$  are given in Table 3. Graph  $(N_e, \nu, \rho_e, t)$  is shown on Figure 3.

$\rho_p(p, x)$	$x = 0$	$x = 1$	$x = 2$	$x = 3$	$x = 4$	$x = 5$	$x = 6$	$x = 7$
$\rho_0(0, x)$	0	0.1	0.3	0.1	0.7	0.1	0.3	0.1
$\rho_1(1, x)$	0.1	0	0.1	0	0.1	0	0.1	0
$\rho_2(2, x)$	0.3	0.1	0	0.1	0.3	0.1	0.3	0.1
$\rho_3(3, x)$	0.1	0	0.1	0	0.1	0	0.1	0
$\rho_4(4, x)$	0.7	0.1	0.3	0.1	0	0.1	0.3	0.1
$\rho_5(5, x)$	0.1	0	0.1	0	0.1	0	0.1	0
$\rho_6(6, x)$	0.3	0.1	0	0.1	0.3	0.1	0.3	0.1
$\rho_7(7, x)$	0.1	0	0.1	0	0.1	0	0.1	0

Table 3:  $\rho_p(p, x)$  when  $\nu_t = \{0, 2, 4, 6\}$ 

PROPOSITION 3.4. Let  $\nu$  be a fuzzy ideal of  $N$  and  $t \in (\alpha, \beta]$ .

- (i) If  $p \in \nu_t$  then  $(N_e, \nu, \rho_p, t)$  is a star graph with root vertex  $p$ .
- (ii) If  $\nu$  is an equiprime fuzzy ideal of  $N$  and  $(N_e, \nu, \rho_p, t)$  is a star graph with root vertex  $p$  then  $p \in \nu_t$ .
- (iii) Let  $x \in N$ . If  $x \in \nu_t$  then  $x$  is connected to all other vertices of  $(N_e, \nu, \rho_e, t)$ .

*Proof.* Let  $t \in (\alpha, \beta]$ . Then  $\nu_t$  is an ideal of  $N$ . To prove (i), let  $p \in \nu_t$  and  $x \in N$ . Let  $r \in N$  be arbitrarily fixed. By the property of ideal,  $prx \in \nu_t$  and  $pr0 \in \nu_t$ . Then  $prx - pr0 \in \nu_t$  (by property of ideal.) Hence  $p$  is adjacent to  $x$  in  $(N_e, \nu, \rho_p, t)$ .




Figure 3:  $(N_e, \nu, \rho_e, t)$  when  $\nu_t = \{0, 2, 4, 6\}$

Therefore  $p$  is adjacent to all other vertices of  $(N_e, \nu, \rho_p, t)$ . Thus  $(N_e, \nu, \rho_p, t)$  is a star graph with root vertex  $p$ .

To prove (ii), let  $(N_e, \nu, \rho_p, t)$  be a star graph with root vertex  $p$ . Then by definition for any  $x$  in  $N$ ,  $prx - pr0 \in \nu_t$  for all  $r \in N$  or  $xrp - xr0 \in \nu_t$  for all  $r \in N$ . Without loss of generality assume  $prx - pr0 \in I$  for all  $r \in N$ . Suppose  $\nu_t = N$ . Then  $p \in \nu_t$ . Suppose  $\nu_t \subset N$ . Choose  $x \in N \setminus \nu_t$ . By the property of equiprime fuzzy ideal of  $N$ , we get  $\nu_t$  is an equiprime ideal of  $N$ . Then we get  $p \in \nu_t$ .

To prove (iii), let  $x \in \nu_t$ . Then by (i),  $(N_e, \nu, \rho_x, t)$  is a star graph with root vertex  $x$ . Then  $x$  is adjacent with all other vertices of  $N$  in  $(N_e, \nu, \rho_x, t)$ . By definition of equiprime fuzzy graph we get  $x$  is adjacent to all other vertices of  $(N_e, \nu, \rho_e, t)$ .  $\square$

Now we provide examples to show the conditions in Proposition 3.4 are necessary.

- (i) In Example 3.2, note that  $a \notin \nu_t$ . Observe that  $(N_e, \nu, \rho_a, t)$  is not a star graph.
- (ii) Now we provide Example 3.5 to show that if  $\nu$  is not an equiprime fuzzy ideal of  $N$  then even if  $(N_e, \nu, \rho_p, t)$  is a star graph with root vertex  $p$  we get  $p \notin \nu_t$ .

EXAMPLE 3.5. Let  $N$  be the nearring with addition and multiplication defined as in Table 1. Define  $\nu : N \rightarrow (0, 1]$  by

$$\nu(x) = \begin{cases} 0.8 & \text{if } x = 0, \\ 0.5 & \text{if } x = b, \\ 0.1 & \text{if } x \in \{a, c\}. \end{cases}$$

If thresholds are  $\alpha = 0.5$  and  $\beta = 0.8$  then  $\nu$  is a fuzzy ideal of  $N$ . Let  $t = \beta$ . Then  $\nu_t = \{0\}$  and the values of  $\rho_b(b, x)$  are given in Table 4. Graph  $(N_e, \nu, \rho_b, t)$  is shown in Figure 4.

	$x = 0$	$x = a$	$x = b$	$x = c$
$\rho_b(b, x)$	0.5	0.1	0	0.1

Table 4:  $\rho_b(b, x)$  when  $\nu_t = \{0\}$




Figure 4:  $(N_e, \nu, \rho_b, t)$  when  $\nu_t = \{0\}$

Note that  $\nu$  is not an equiprime fuzzy ideal of  $N$  ( $\alpha \vee \mu(b) \vee \mu(b-0) = 0.5 \vee 0.5 \vee 0.5 = 0.5 \not\geq 0.8 = 0.8 \wedge 0.8 = \beta \wedge \inf_{r \in N} \mu(brb - br0)$ ) and  $(N_e, \nu, \rho_b, t)$  is a star graph with root vertex  $b$ . Observe that  $b \notin \nu_t$ .

(iii) In Example 3.3, note that  $7 \notin \nu_t$ . Observe that 7 is not connected to all other vertices in  $(N_e, \nu, \rho_e, t)$ .

**PROPOSITION 3.6.** *Let  $\nu$  be a fuzzy ideal of  $N$  and  $t \in (\alpha, \beta]$ .*

- (i) *Let  $\nu$  be an equiprime fuzzy ideal of  $N$ . Then  $\nu_t$  is a strong vertex cut of  $(N_e, \nu, \rho_e, t)$ .*  
(ii) *Let  $\nu$  be an equisemiprime fuzzy ideal of  $N$  and  $\nu_t$  be a strong vertex cut of  $(N_e, \nu, \rho_e, t)$ . Then  $\nu$  is an equiprime fuzzy ideal of  $N$ .*

*Proof.* To prove (i), let  $\nu$  be an equiprime fuzzy ideal of  $N$  and  $t \in (\alpha, \beta]$ . Suppose  $\nu_t = N$ . Then  $\nu_t$  is a strong vertex cut of  $(N_e, \nu, \rho_e, t)$ . Let  $\nu_t \subset N$ . Let  $p \in N \setminus \nu_t$  and  $x \in N \setminus \nu_t$  such that there exists an edge between  $p$  and  $x$  in  $(N_e, \nu, \rho_e, t)$ . Then  $prx - pr0 \in \nu_t$  or  $xrp - xr0 \in \nu_t$  for all  $r \in N$ . Without loss of generality assume  $prx - pr0 \in \nu_t$  for all  $r \in N$ . As  $\nu$  is an equiprime fuzzy ideal of  $N$ , we get  $\nu_t$  is an equiprime ideal of  $N$ . Then  $p \in \nu_t$  or  $x \in \nu_t$ . A contradiction to the fact that  $p \in N \setminus \nu_t$  and  $x \in N \setminus \nu_t$ . Hence  $\nu_t$  is a strong vertex cut of  $(N_e, \nu, \rho_e, t)$ .

To prove (ii), let  $\nu$  be an equisemiprime fuzzy ideal of  $N$  and  $\nu_t$  be a strong vertex cut of  $(N_e, \nu, \rho_e, t)$ . Let  $prx - pro \in \nu_t$  for all  $r \in N$ . Suppose  $p = x$ . Then  $p \in \nu_t$  ( $\nu_t$  is equisemiprime.) Let  $p \neq x$ . Suppose  $p \in N \setminus \nu_t$  and  $x \in N \setminus \nu_t$ . As  $\nu_t$  is a strong vertex cut of  $(N_e, \nu, \rho_e, t)$  there is no edge between  $p$  and  $x$  in  $(N_e, \nu, \rho_e, t)$ . Then  $prx - pro \notin \nu_t$  and  $xrp - xr0 \notin \nu_t$  for some  $r \in N$ . A contradiction since  $prx - pro \in \nu_t$  for all  $r \in N$ . Hence  $\nu_t$  is an equiprime ideal of  $N$ . Therefore  $\nu$  is an equiprime fuzzy ideal of  $N$ .  $\square$

Now we provide examples to show the conditions in Proposition 3.6 are necessary.

- (i) We provide Example 3.7 to show that if  $\nu$  is not an equiprime fuzzy ideal of  $N$ . Then  $\nu_t$  not a strong vertex cut of  $(N_e, \nu, \rho_e, t)$ .  
(ii) We provide Example 3.8 to show that if  $\nu$  is not an equisemiprime fuzzy ideal of  $N$ . Then even if  $\nu_t$  is a strong vertex cut of  $(N_e, \nu, \rho_e, t)$ ,  $\nu$  is not an equiprime fuzzy ideal of  $N$ .

**EXAMPLE 3.7.** Let  $N$  be the nearring with addition  $+$  and multiplication  $\cdot$  defined as in Table 1. We define  $\nu : N \rightarrow (0, 1]$  by

$$\nu(x) = \begin{cases} 0.8 & \text{if } x = 0, \\ 0.5 & \text{if } x = b, \\ 0.1 & \text{if } x \in \{a, c\}. \end{cases}$$

If we take thresholds  $\alpha = 0.5$  and  $\beta = 0.8$  then  $\nu$  is a fuzzy ideal of  $N$ . Let  $t = \beta$ . Then  $\nu_t = \{0\}$  and for  $p \in N$  the values of  $\rho_p(p, x)$  are given in Table 5. Graph  $(N_e, \nu, \rho_e, t)$  is shown in Figure 5.



$\rho_p(p, x)$	$x = 0$	$x = a$	$x = b$	$x = c$
$\rho_0(0, x)$	0	0.1	0.1	0.1
$\rho_a(a, x)$	0.1	0	0.5	0
$\rho_b(b, x)$	0.5	0.1	0	0.1
$\rho_c(c, x)$	0.1	0	0.1	0

Table 5:  $\rho_0(x, y)$  when  $\nu_t = \{0\}$




Figure 5:  $(N_e, \nu, \rho_e, t)$  when  $\nu_t = \{0\}$

Note that  $\nu$  is not an equiprime fuzzy ideal of  $N$  ( $\alpha \vee \nu(b) \vee \nu(b-0) = 0.5 \vee 0.5 \vee 0.5 = 0.5 \not\geq 0.8 = 0.8 \wedge 0.8 = \beta \wedge \inf_{r \in N} \nu(brb - br0)$ .) Observe that  $\nu_t = \{0\}$  is not a strong vertex cut of  $(N_e, \nu, \rho_e, t)$ .

EXAMPLE 3.8. Let  $N = \mathbb{Z}_4$  be the ring of integers modulo 4. We define  $\nu : N \rightarrow [0, 1]$  by

$$\nu(x) = \begin{cases} 0.9 & \text{if } x = 0, \\ 0.6 & \text{if } x = 2, \\ 0.3 & \text{if } x \in \{1, 3\}. \end{cases}$$

Take thresholds  $\alpha = 0.6$  and  $\beta = 0.9$ . Then  $\nu$  is a fuzzy ideal of  $N$ . Let  $t = \beta$ . Then  $\nu_t = \{0\}$  and the values of  $\rho_p(p, x)$  are as in Table 6. The graph is given in Figure 6.

$\rho_p(p, x)$	$x = 0$	$x = 1$	$x = 2$	$x = 3$
$\rho_0(0, x)$	0	0.3	0.6	0.3
$\rho_1(1, x)$	0.3	0	0	0
$\rho_2(2, x)$	0.6	0	0	0
$\rho_3(3, x)$	0.3	0	0	0

Table 6: Values of  $\rho_p(p, x)$  when  $\nu_t = \{0\}$




Figure 6:  $(N_e, \nu, \rho_e, t)$  when  $\nu_t = \{0\}$

Note that  $\nu$  is not an equisemiprime fuzzy ideal of  $N$  since  $\alpha \vee \nu(2) \vee \nu(2) = 0.6 \vee 0.6 \vee 0.6 = 0.6 \not\geq 0.9 = 0.9 \wedge 0.9 = \beta \wedge \inf_{r \in N} \nu(2r2 - 2r0)$  and  $\nu_t = \{0\}$  is a strong vertex cut of  $(N_e, \nu, \rho_e, t)$ . Observe that  $\nu$  is not an equiprime fuzzy ideal of  $N$ .

PROPOSITION 3.9. Let  $\nu$  be an equiprime fuzzy ideal of  $N$  and  $t \in (\alpha, \beta]$ .

(i) Let  $x$  be a vertex in  $(N_e, \nu, \rho_e, t)$ . If  $\rho_p(p, x) > 0 \forall p \neq x; p \in N$  then  $p \in \nu_t$ .

(ii) Let  $x$  be a vertex in  $(N_e, \nu, \rho_e, t)$ . If  $p \in \nu_t$  then  $\rho_p(p, x) > 0 \forall p \neq x; p \in N$ .

(iii)  $\nu$  is an equiprime fuzzy ideal of  $N$  if and only if every element  $x \in \nu_t$  is connected to all other elements of  $N$  in  $(N, \nu, \rho_e, t)$ .

(iv) If  $\nu$  is an equiprime fuzzy ideal  $N$  then  $(\nu_t, \nu, \rho, t)$  is a complete subgraph of  $(N, \nu, \rho_e, t)$ .

REMARK 3.10. (i) In Example 3.2, note that  $\nu$  is an equiprime fuzzy ideal of  $N$  and  $\rho_a(a, c) = 0$  for  $x = c$ . Observe that  $a \notin \nu_t$ .

(ii) In Example 3.2, note that  $c \notin \nu_t$ . Observe that  $\rho_c(c, a) = 0$ .

DEFINITION 3.11. Let  $(N_e, \nu, \rho_e, t)$  be an equiprime fuzzy graph. Then  $(N_e, \nu, \rho_e, t)$  is said to be *ideal symmetric* if for every pair of vertices  $a, b$  in  $(N_e, \nu, \rho_e, t)$  with an edge between them, either  $[\rho(a, c) > 0 \forall c \neq a; c \in N]$  or  $[\rho(b, c) > 0 \forall c \neq b; c \in N]$ .

PROPOSITION 3.12. Let  $\nu$  be a fuzzy ideal of  $N$  and  $t \in (\alpha, \beta]$ .

(a) If  $\nu$  is an equiprime fuzzy ideal of  $N$  then  $(N_e, \nu, \rho_e, t)$  is ideal symmetric.

(b) Suppose (i)  $(N, \nu, \rho_e, t)$  is ideal symmetric; (ii)  $\nu$  is equisemiprime fuzzy ideal of  $N$ ; (iii) for every  $x \in N$  and  $\rho(x, z) > 0 \forall z \neq x; z \in N$  implies  $x \in \nu_t$ . Then  $\nu$  is an equiprime fuzzy ideal of  $N$ .

*Proof.* Let  $t \in (\alpha, \beta]$ . To prove (a) let  $\nu$  be an equiprime fuzzy ideal of  $N$  and  $p, x \in N$  be such that there is an edge between  $p$  and  $x$  in  $(N_e, \nu, \rho_e, t)$ . Then for all  $r \in N$  we get  $prx - pr0 \in \nu_t$  or  $xrp - xr0 \in \nu_t$ . Without loss of generality, assume  $prx - pr0 \in \nu_t$ . Then  $p \in \nu_t$  or  $x \in \nu_t$  (as  $\nu$  is an equiprime fuzzy ideal of  $N$  then  $\nu_t$  is an equiprime ideal of  $N$ ). By Proposition 3.9 (ii) we get  $\rho(p, z) > 0 \forall z \neq p; z \in N$  or  $\rho(x, z) > 0 \forall z \neq x; z \in N$ . Hence  $(N, \nu, \rho, t)$  is ideal symmetric.

To prove (b), let  $p, x \in N$  and  $prx - pr0 \in \nu_t$  for all  $r \in R$ . Suppose  $\nu_t = N$ . Then  $p \in \nu_t$ . Let  $\nu_t \subset N$ . Suppose  $p = x$ . Then  $p \in \nu_t$  (by (ii)  $\nu$  is an equisemiprime fuzzy ideal of  $N$  and  $\nu_t$  is an equisemiprime ideal of  $N$ ). Let  $p \neq x$ . Now there exists an edge between  $p$  and  $x$  in  $(N_e, \nu, \rho, t)$ . As  $(N_e, \nu, \rho, t)$  is ideal symmetric we get either  $\rho(p, z) > 0 \forall z \neq p; z \in N$  or  $\rho(x, z) > 0 \forall z \neq x; z \in N$ . By Proposition 3.9 (i), we get  $p \in \nu_t$  or  $x \in \nu_t$ . Thus  $\nu$  is an equiprime fuzzy ideal of  $N$ .  $\square$

If  $\nu$  is not an equiprime fuzzy ideal of  $N$  then  $(N_e, \nu, \rho_e, t)$  is not ideal symmetric. We provide Example 3.13.

EXAMPLE 3.13. Let  $N = \mathbb{Z}_6$  be the ring of integers modulo 6. We define  $\nu : N \rightarrow (0, 1]$  by

$$\nu(x) = \begin{cases} 0.9 & \text{if } x = 0, \\ 0.7 & \text{if } x \in \{2, 4\}, \\ 0.3 & \text{if } x \in \{1, 3, 5\}. \end{cases}$$

If we take thresholds  $\alpha = 0.7$  and  $\beta = 0.9$  then  $\nu$  is a fuzzy ideal of  $N$ . Let  $t = \beta$ . Then  $\nu_t = \{0\}$  and the values of  $\rho_p(p, x)$  are given in Table 7. The graph  $(N_e, \nu, \rho_e, t)$  is shown in Figure 7.

$\rho_p(p, x)$	$x = 0$	$x = 1$	$x = 2$	$x = 3$	$x = 4$	$x = 5$
$\rho_0(0, x)$	0	0.5	0.7	0.5	0.7	0.5
$\rho_1(1, x)$	0.5	0	0	0	0	0
$\rho_2(2, x)$	0.7	0	0	0.5	0	0
$\rho_3(3, x)$	0.5	0	0.5	0	0.5	0
$\rho_4(4, x)$	0.7	0	0	0.5	0	0
$\rho_5(5, x)$	0.5	0	0	0	0	0

Table 7:  $\rho_p(p, x)$  when  $\nu_t = \{0\}$




Figure 7:  $(N_e, \nu, \rho_e, t)$  when  $\nu_t = \{0\}$

Note that  $\nu$  is not an equirpeime fuzzy ideal of  $N$ .  $(\alpha \vee \nu(2) \vee \nu(3) = 0.7 \vee 0.7 \vee 0.3 = 0.7 \not\geq 0.9 = 0.9 \wedge 0.9 = \beta \wedge \inf_{r \in N} \nu(2r3 - 2r0))$ . Observe that the graph  $(N_e, \nu, \rho_e, t)$  in Figure 7 is not ideal symmetric.

In Example 3.8, note that the graph in Figure 6 is ideal symmetric, however  $\nu$  is not an equisemiprime fuzzy ideal of  $N$ . Since  $\alpha \vee \nu(2) \vee \nu(2) = 0.6 \vee 0.6 \vee 0.6 = 0.6 \not\geq 0.9 = 0.9 \wedge 0.9 = \beta \wedge \inf_{r \in N} \nu(2r2 - 2r0)$ . Observe that  $\nu$  is not an equiprime fuzzy ideal of  $N$ .

PROPOSITION 3.14. *Let  $\nu$  be an equiprime fuzzy ideal of  $N$ ,  $t \in (\alpha, \beta]$  and  $n \notin \nu_t$ . Then  $C = \nu_t \cup \{n\}$  is a fuzzy clique of  $(N_e, \nu, \rho_e, t)$ .*

*Proof.* Let  $t \in (\alpha, \beta]$ . Then by Proposition 3.9 (iv), we get  $(\nu_t, \nu, \rho, t)$  is complete subgraph of  $(N_e, \nu, \rho_e, t)$ . Let  $n \notin \nu_t$  and  $C = \nu_t \cup \{n\}$ . By Proposition 3.9 (iii), we get that every element of  $\nu_t$  is connected to  $n$ . This proves that  $(C, \nu, \rho_e, t)$  is complete. It remains to prove that  $C$  is maximal. First, we show that if  $p \in N \setminus \nu_t, x \in N \setminus \nu_t$  and  $p \neq x$  then  $p$  is not connected to  $x$  in  $(N_e, \nu, \rho_e, t)$ . If possible, let  $p$  be connected to  $x$ . Then for all  $r \in N$  we get  $prx - pr0 \in \nu_t$  or  $xrp - xr0 \in \nu_t$ . Without loss of generality, assume  $prx - pr0 \in \nu_t$ . As  $\nu_t$  is an equiprime ideal of  $N$ , we get  $p \in \nu_t$  or  $x \in \nu_t$ . This is a contradiction to the fact that  $p \in N \setminus \nu_t, x \in N \setminus \nu_t$ . Now, let  $C_1 = C \cup \{m\}$ ;  $m \notin \nu_t$  and  $m \neq n$ . Then  $\rho_e(n, m) = 0 \neq \nu(m) \wedge \nu(n)$ . This implies  $(C_1, \nu, \rho_e, t)$  is not complete. Hence  $(C_1, \nu, \rho_e, t)$  cannot be a fuzzy clique. Thus  $C = \nu_t \cup \{n\}$  is a fuzzy clique of  $(N_e, \nu, \rho_e, t)$ .  $\square$




Figure 8: Complete subgraphs of  $(N_e, \nu, \rho_e, t)$

REMARK 3.15. (i) In Example 3.2, take  $b \notin \nu_t$ . Then observe that  $C = \nu_t \cup \{b\}$  is a fuzzy clique of  $(N_e, \nu, \rho_e, t)$  as shown in Figure 8a.

(ii) In Example 3.3, take  $3 \notin \nu_t$ . Then observe that  $C = \nu_t \cup \{3\}$  is a fuzzy clique of  $(N_e, \nu, \rho_e, t)$  as shown in Figure 8b.

(iii) In Example 3.13, note that  $\nu$  is not an equiprime fuzzy ideal of  $N$ . If we take  $3 \notin \nu_t$  then observe that  $C = \nu_t \cup \{3\}$  is not a fuzzy clique of  $(N_e, \nu, \rho_e, t)$  as shown in Figure 8c.

PROPOSITION 3.16. *Let  $\nu$  be an equiprime fuzzy ideal of  $N$  and  $t \in (\alpha, \beta]$ . Then*

(i)  $\nu_t$  is a vertex cover of  $(N_e, \nu, \rho_e, t)$ . (ii)  $(N_e \setminus \nu_t, \nu, \rho_e, t)$  is an empty graph.

REMARK 3.17. (i) In Example 3.2, note that  $\nu$  is an equiprime fuzzy ideal of  $N$ . Observe that  $\nu_t$  is a vertex cover of  $(N_e, \nu, \rho_e, t)$ .

(ii) In Example 3.13, note that  $\nu$  is not an equiprime fuzzy ideal of  $N$ . Note that  $\nu_t = \{0\}$  is not a vertex cover of  $(N_e, \nu, \rho_e, t)$ .

(iii) In Example 3.13, note that  $\nu$  is not an equiprime fuzzy ideal of  $N$ . Note that  $(N_e \setminus \nu_t, \nu, \rho_e, t)$  is not an empty graph. In fact  $(N_e \setminus \nu_t, \nu, \rho_e, t)$  is as shown in Figure 9.




Figure 9:  $(N_e, \nu, \rho_e, t)$  when  $\nu_t = \{0\}$

PROPOSITION 3.18. *Let  $N$  be a simple nearring and  $\nu$  be an equiprime fuzzy ideal of  $N$ . Then  $(N_e, \nu, \rho_e, t)$  is a star graph or  $(N_e, \nu, \rho_e, t)$  is a complete graph.*

REMARK 3.19. In Example 3.2, note that  $\nu$  is an equiprime fuzzy ideal, however the nearring is not a simple nearring. Observe that the graph is neither a star graph nor a complete graph.

#### 4. Interrrelations between fuzzy graphs

PROPOSITION 4.1. *Let  $\nu$  be a fuzzy ideal of  $N$  and  $t \in (\alpha, \beta]$ . Then  $(N_3, \nu, \rho_3, t) = (N_e, \nu, \rho_e, t)$  if any one of the following conditions are satisfied.*

- (i)  $N$  is zero symmetric. (ii)  $N$  is distributive.  
 (iii) Every ideal of  $N$  is totally reflexive. (iv)  $N$  is equiprime.

*Proof.* Let  $t \in (\alpha, \beta]$ . Then  $\nu_t$  is an ideal of  $N$  for all  $t \in (\alpha, \beta]$ . We have vertex set of  $(N_3, \nu, \rho_3, t) = (N_e, \nu, \rho_e, t) = N$ . To prove (i), suppose  $p, x \in N$  such that there is an edge between  $p$  and  $x$  in  $(N_3, \nu, \rho_3, t)$ . Then  $pNx \subseteq \nu_t$  or  $xNp \subseteq \nu_t$ . Without loss of generality assume  $pNx \subseteq \nu_t$ . Then  $pnx \in \nu_t$  for all  $n \in N$ . Let  $n \in N$  is arbitrarily fixed. Then we get  $pnx - pn0 \in \nu_t$ . (as  $N$  is zero symmetric  $xn0 = 0$ ) Then  $p$  and  $x$  are

adjacent in  $(N_e, \nu, \rho_e, t)$ . Therefore  $E((N_3, \nu, \rho_3, t)) \subseteq E((N_e, \nu, \rho_e, t))$ . Now suppose  $p, x \in N$  such that  $(p, x)$  is an edge in  $(N_e, \nu, \rho_e, t)$ . Then  $prx - pr0 \in \nu_t$  for all  $r \in N$  or  $xnp - xn0 \in \nu_t$  for all  $r \in N$ . Without loss of generality assume  $prx - pr0 \in \nu_t$  for all  $r \in N$ . Let  $r \in N$  be arbitrarily fixed. Then we get  $prx \in N$  (as  $N$  is zero symmetric we get  $pr0 = 0$ ). Hence we get  $prx \in \nu_t$  for all  $r \in N$ . Then  $pNx \subseteq \nu_t$ , which implies  $p$  and  $x$  are adjacent in  $(N_3, \nu, \rho_3, t)$ . Hence  $E((N_e, \nu, \rho_e, t)) \subseteq E((N_3, \nu, \rho_3, t))$ . Therefore  $E((N_e, \nu, \rho_e, t)) = E((N_3, \nu, \rho_3, t))$ . Thus  $(N_3, \nu, \rho, t) = (N_e, \nu, \rho_e, t)$ .

The proofs of (ii), (iii) and (iv) are similar to that of (i).  $\square$

We provide examples to show that if conditions given in Proposition 4.1 are not satisfied then  $(N_3, \nu, \rho_3, t) \neq (N_e, \nu, \rho_e, t)$ .

EXAMPLE 4.2. Let  $N = \{0, a, b, c\}$  be the nearring with addition and multiplication defined as in Table 1. We define  $\nu : N \rightarrow (0, 1]$  by

$$\nu(x) = \begin{cases} 0.9 & \text{if } x = 0, \\ 0.4 & \text{if } x = b, \\ 0.2 & \text{if } x \in \{a, c\}. \end{cases}$$

If we take thresholds  $\alpha = 0.4$  and  $\beta = 0.9$  then  $\nu$  is a fuzzy ideal of  $N$ . Let  $t = \beta$ . Then  $\nu_t = \{0\}$  and the values of  $\rho_p(p, x)$  are as in Table 8 and the graph  $(N_e, \nu, \rho_e, t)$  is given in Figure 10.

$\rho_p(p, x)$	$x = 0$	$x = a$	$x = b$	$x = c$
$\rho_0(0, x)$	0	0.2	0.4	0.1
$\rho_a(a, x)$	0.2	0	0.4	0
$\rho_b(b, x)$	0.4	0.2	0	0.2
$\rho_c(c, x)$	0.2	0	0.2	0

Table 8: Values of  $\rho_p(p, x)$  when  $\nu_t = \{0\}$




Figure 10:  $(N_e, \nu, \rho, t)$  for  $\nu_t = \{0\}$

The values of  $\rho(x, y)$  are as in Table 9 and the graph  $(N_3, \nu, \rho_3, t)$  is given in Figure 11.

$\rho(x, y)$	$y = 0$	$y = a$	$y = b$	$y = c$
$x = 0$	0	0.2	0.4	0.1
$x = a$	0.2	0	0.4	0
$x = b$	0.4	0.4	0	0
$x = c$	0.2	0	0	0

Table 9: Values of  $\rho(x, y)$  when  $\nu_t = \{0\}$




Figure 11:  $(N_3, \nu, \rho_3, t)$  for  $\nu_t = \{0\}$

In this example, note that:

- (i) The nearring  $N$  is not zero symmetric ( $b \cdot 0 \neq 0, c \cdot 0 \neq 0$ ). Observe that  $(N_3, \nu, \rho, t) \neq (N_e, \nu, \rho, t)$ .
- (ii) The nearring  $N$  is not distributive ( $b(a + c) = b \cdot b = b$  and  $b \cdot a + b \cdot c = b + b = 0$ ). Hence  $b(a + c) \neq b \cdot a + b \cdot c$ . Observe that  $(N_3, \nu, \rho, t) \neq (N_e, \nu, \rho, t)$ .
- (iii) The ideal  $\nu_t$  is not totally reflexive ( $aNb = \{0\} \in \nu_t$  however  $bNa = \{b\} \notin \nu_t$ ). Observe that  $(N_3, \nu, \rho_3, t) \neq (N_e, \nu, \rho_e, t)$ .

(iv)  $\{0\}$  is not an equiprime ideal of  $N$ . Hence  $N$  is not an equiprime nearring. Observe that  $(N_3, \nu, \rho_3, t) \neq (N_e, \nu, \rho_e, t)$ .

PROPOSITION 4.3. *Let  $\nu$  be a fuzzy ideal of  $N$  and  $t \in (\alpha, \beta]$ .*

(i) *If  $\nu$  is a 3-prime fuzzy ideal of  $N$  then  $(N_3, \nu, \rho_3, t) \subseteq (N_c, \nu, \rho_c, t)$ .*

(ii) *If  $\nu$  is a c-prime fuzzy ideal of  $N$  then  $(N_3, \nu, \rho_3, t) = (N_c, \nu, \rho_c, t)$ .*

*Proof.* We have  $V((N_3, \nu, \rho, t)) = V((N_c, \nu, \rho, t)) = N$ . To prove (i), let  $\nu$  be a 3-prime fuzzy ideal of  $N$ . Let  $x, y \in N$  such that  $(x, y) \in E((N_3, \nu, \rho_3, t))$ . Then  $xNy \subseteq \nu_t$  or  $yNx \subseteq \nu_t$ . Without loss of generality assume  $xNy \subseteq \nu_t$ . As  $\nu$  is a 3-prime fuzzy ideal of  $N$  we get  $\nu_t$  is a 3-prime ideal of  $N$ . Then we get  $x \in \nu_t$  or  $y \in \nu_t$ . Now, let us assume  $x \in \nu_t$ . Then  $xy \in \nu_t$  for all  $y \in N$ . Then  $(x, y) \in E((N_c, \nu, \rho_c, t))$ . Proof is similar for  $y \in \nu_t$ . Hence  $E((N_3, \nu, \rho_3, t)) \subseteq E((N_c, \nu, \rho_c, t))$ . Therefore  $(N_3, \nu, \rho_3, t) \subseteq (N_c, \nu, \rho_c, t)$ .

To prove (ii), let  $\nu$  be a c-prime fuzzy ideal of  $N$ . Let  $a, b \in N$  such that  $(a, b) \in E((N_c, \nu, \rho_c, t))$ . Then  $ab \in \nu_t$  or  $ba \in \nu_t$ . Without loss of generality assume  $ab \in \nu_t$ . As  $\nu$  is a c-prime fuzzy ideal of  $N$  we get  $\nu_t$  is a c-prime ideal of  $N$ . Then we get  $x \in \nu_t$  or  $y \in \nu_t$ . Now, let us assume  $x \in \nu_t$ . Then by the property of ideal we get  $xN \subseteq \nu_t$ . Then  $xNy \subseteq \nu_t$  for all  $y \in N$ . Then  $(x, y) \in E((N_3, \nu, \rho_3, t))$ . Proof is similar for  $y \in \nu_t$ . Hence  $E((N_c, \nu, \rho_c, t)) \subseteq E((N_3, \nu, \rho_3, t))$ . Therefore  $(N_c, \nu, \rho_c, t) \subseteq (N_3, \nu, \rho_3, t)$ . Every c-prime fuzzy ideal is a 3-prime fuzzy ideal. Then from (i) we get  $E((N_3, \nu, \rho_3, t)) \subseteq E((N_c, \nu, \rho_c, t))$ . Therefore  $(N_3, \nu, \rho_3, t) = (N_c, \nu, \rho_c, t)$ .  $\square$

PROPOSITION 4.4. *Let  $\nu$  be a fuzzy ideal of  $N$  and  $t \in (\alpha, \beta]$ . Then  $(N_3, \nu, \rho_3, t) = (N_c, \nu, \rho_c, t) = (N_e, \nu, \rho_e, t)$ , if any one of the following conditions are satisfied.*

(i)  *$\nu$  is a c-prime fuzzy ideal of a zero symmetric nearring  $N$ .*

(ii)  *$\nu$  is a prime fuzzy ideal of a commutative ring  $N$ .*

## 5. Nearing homomorphism and graph homomorphism

PROPOSITION 5.1. *Let  $f : N_1 \rightarrow N_2$  be an one to one and onto nearring homomorphism. Let  $\nu$  be a fuzzy ideal of  $N_1$  and  $t \in (\alpha, \beta]$ . Then  $f$  is an one to one and onto graph homomorphism from  $(N_{1e}, \nu, \rho_e, t)$  to  $(N_{2e}, f(\nu), f(\rho)_e, t)$ .*

*Proof.* Let  $\nu$  be a fuzzy ideal of  $N_1$  and  $t \in (\alpha, \beta]$ . Then by [18, Proposition 3.25] we get  $f(\nu)$  is a fuzzy ideal of  $N_2$  with same thresholds as that of  $\nu$ . Let  $p, x \in N_1$  such that  $p \neq x$  and  $(p, x)$  be an edge of  $(N_{1e}, \nu, \rho_e, t)$ . Then for all  $r \in N_1$  we get  $prx - pr0_1 \in \nu_t$  or  $xrp - xr0_1 \in \nu_t$  where  $0_1$  is the additive identity of  $N_1$ . Without loss of generality assume  $prx - pr0_1 \in \nu_t$ . Let  $r \in N_1$  be arbitrarily fixed. Then  $f(prx - pr0_1) \in f(\nu_t) \subseteq f(\nu)_t$  (by [18, Remark 3.30]). Then  $f(prx - pr0_1) \in f(\nu)_t$ . As  $f$  is a nearring homomorphism we get  $f(prx - pr0_1) = f(p)f(r)f(x) - f(p)f(r)f(0_1) = f(p)f(r)f(x) - f(p)f(r)0_2 \in f(\nu)_t$  where  $0_2$  is the additive identity of  $N_2$ . As  $f$  is one to one we get  $f(p) \neq f(x)$ . Also  $f(N_1) = N_2$  ( $f$  is onto). Hence  $(f(p), f(x))$  is an edge in  $(N_{2e}, f(\nu), f(\rho)_e, t)$ . Therefore  $f$  is a graph homomorphism

from  $(N_{1e}, \nu, \rho_e, t)$  to  $(N_{2e}, f(\nu), f(\rho)_e, t)$ . Let  $p_1, p_2 \in N_1$  such that  $p_1 \neq p_2$ . Then  $f(p_1) \neq f(p_2)$  (since  $f$  is one to one). Hence there is one to one correspondence between the vertex set of  $(N_{1e}, \nu, \rho_e, t)$  and  $(N_{2e}, f(\nu), f(\rho)_e, t)$ . Let  $(p_1, x_1)$  and  $(p_2, x_2)$  be edges of  $(N_{1e}, \nu, \rho_e, t)$  such that  $(p_1, x_1) \neq (p_2, x_2)$ . Then for all  $r_1 \in N_1$  we get  $(p_1 r_1 x_1 - p_1 r_1 0_1) \in \nu_t$  or  $(x_1 r_1 p_1 - x_1 r_1 0_1) \in \nu_t$  and for all  $r_2 \in N_1$  we get  $(p_2 r_2 x_2 - p_2 r_2 0_1) \in \nu_t$  or  $(x_2 r_2 p_2 - x_2 r_2 0_1) \in \nu_t$ . Without loss of generality assume  $(p_1 r_1 x_1 - p_1 r_1 0_1) \in \nu_t$  and  $(p_2 r_2 x_2 - p_2 r_2 0_1) \in \nu_t$ . Let  $r_1, r_2 \in N_1$  be arbitrarily fixed. Then  $f(p_1 r_1 x_1 - p_1 r_1 0_1) = f(p_1) f(r_1) f(x_1) - f(p_1) f(r_1) f(0_1) \in f(\nu)_t$  and  $f(p_2 r_2 x_2 - p_2 r_2 0_1) = f(p_2) f(r_2) f(x_2) - f(p_2) f(r_2) f(0_2) \in f(\nu)_t$ . As  $f$  is one to one we get  $f(p_1) f(r_1) f(x_1) - f(p_1) f(r_1) f(0_1) \neq f(p_2) f(r_2) f(x_2) - f(p_2) f(r_2) f(0_2)$ . Therefore there is one to one correspondence between the edge sets of  $(N_{1e}, \nu, \rho_e, t)$  and  $(N_{2e}, f(\nu), f(\rho)_e, t)$ .  $\square$

EXAMPLE 5.2. Let  $Z_n$  be the ring of integers modulo  $n$ . let  $N$  be the nearring defined in Example 3.2. Let  $N_1 = Z_1 \times N = \{(0, 0), (0, a), (0, b), (0, c)\}$  and  $N_2 = N$ . Define  $f : N_1 \rightarrow N_2$  by  $f((x, y)) = y$ . Then  $f$  is an one to one and onto nearring homomorphism. We define  $\nu : N_1 \rightarrow (0, 1]$  by

$$\nu(x) = \begin{cases} 0.9 & \text{if } x = (0, 0), \\ 0.5 & \text{if } x = (0, b), \\ 0.1 & \text{if } x \in \{(0, a), (0, c)\}. \end{cases}$$

If we take thresholds  $\alpha=0.1$  and  $\beta=0.5$  then  $\nu$  is a fuzzy ideal of  $N_1$ . Let  $t=\beta$ . Then  $\nu_t = \{(0, 0), (0, b)\}$  and the values of  $\rho_p(p, x)$  are as in Table 10 and the graph is given in Figure 12.

$\rho_p(p, x)$	$x = (0, 0)$	$x = (0, a)$	$x = (0, b)$	$x = (0, c)$
$\rho_{(0,0)}((0, 0), x)$	0	0.1	0.5	0.1
$\rho_{(0,a)}((0, a), x)$	0.1	0	0.5	0
$\rho_{(0,b)}((0, b), x)$	0.5	0.1	0	0.1
$\rho_{(0,c)}((0, c), x)$	0.1	0	0.1	0

Table 10: Values of  $\rho(x, y)$  when  $\nu_t = \{(0, 0), (0, b)\}$




Figure 12:  $(N_e, \nu, \rho_e, t)$  for  $\nu_t = \{(0, 0), (0, b)\}$

Then  $f(\nu)$  is a fuzzy ideal of  $N_2$  with the same thresholds as that of  $\nu$ . Let  $f(\nu) : N_2 \rightarrow (0, 1]$  be given by

$$\nu(x) = \begin{cases} 0.7 & \text{if } x = 0, \\ 0.4 & \text{if } x = b, \\ 0.1 & \text{if } x \in \{a, c\}. \end{cases}$$

Then  $f(\nu_t) = f(\{(0,0), (0,b)\}) = \{0, b\}$  is an ideal of  $N_2$ . The values of  $\rho$  are as in Table 11 and the graph is given in Figure 13.

$\rho_p(p, x)$	$x = 0$	$x = a$	$x = b$	$x = c$
$\rho_0(0, x)$	0	0.1	0.4	0.1
$\rho_a(a, x)$	0.1	0	0.4	0
$\rho_b(b, x)$	0.4	0.1	0	0.1
$\rho_c(c, x)$	0.1	0	0.1	0




Table 11: Values of  $\rho_p(p, x)$  when  $\nu_t = \{0, b\}$  Figure 13:  $(N_e, \nu, \rho_e, t)$  for  $\nu_t = \{0, b\}$  Observe that  $f$  is an one to one graph homomorphism from  $N_1$  to  $N_2$ .

**PROPOSITION 5.3.** *Let  $f : N_1 \rightarrow N_2$  be an one to one and onto nearring homomorphism. Let  $\nu$  be a fuzzy ideal of  $N_1$  and  $t \in (\alpha, \beta]$ . If  $x \in \nu_t$  then  $\rho(f(x), f(z)) > 0 \forall f(z) \neq f(x); f(z) \in N_2$ .*

*Proof.* Let  $t \in (\alpha, \beta]$  and  $x \in \nu_t$ .

**Case 1.** Suppose  $p = 0_1$  where  $0_1$  is the additive identity of  $N_1$ . Then  $f(p) = f(0_1) = 0_2$  where  $0_2$  is the additive identity of  $N_2$ . Then  $(prx - pr0_1) = 0_1 \in \nu_t$  for all  $x \in N_1$ . Let  $r \in N_1$  be arbitrarily fixed. As  $f$  is a nearring homomorphism we get  $f(prx - pr0_1) = f(p)f(r)f(x) - f(p)f(r)f(0_1) = 0_2f(r)f(x) - 0_2f(r)0_2 = 0_2$ , which is true for all  $f(x) \in N_2$ . Then  $\rho(f(p), f(x)) > 0$  for all  $f(p) \neq f(x)$ .

**Case 2.** Let  $p \neq 0_1$ . As  $p \in \nu_t$  and  $\nu_t$  is an ideal, then for  $r \in N_1$  we get  $prx \in \nu_t$  and  $pr0 \in \nu_t$  for all  $x \in N_1$ . Then  $prx - pr0 \in \nu_t$ . Then  $f(prx - pr0) \in f(\nu_t) \subseteq f(\nu)_t$  for all  $f(x) \in N_2$ . As  $f$  is a nearring homomorphism we get  $f(p)f(r)f(x) - f(p)f(r)f(0_1) \in f(\nu)_t$ . As  $f$  is one to one we get  $f(p) \neq f(x)$ . Hence  $(f(p), f(x)) \in E(((N_2)_e, f(\nu), f(\rho)_e, t))$ . Therefore  $\rho(f(p), f(x)) > 0 \forall f(p) \neq f(x); f(x) \in N_2$ .  $\square$

**REMARK 5.4.** In Example 5.2, note that  $(0, a) \notin \nu_t$ . Then  $f((0, a)) = a$  is not connected to all other vertices of  $N_2$ . Also  $(0, b) \in \nu_t$ . Then  $f((0, b)) = b$  is connected to all other vertices of  $N_2$ .

**PROPOSITION 5.5.** *Let  $f : N_1 \rightarrow N_2$  be an one to one and onto nearring homomorphism. Let  $\nu$  be a fuzzy ideal of  $N_1$  and  $t \in (\alpha, \beta]$ . If  $\rho_{f(p)}(f(p), f(x)) > 0 \forall f(p) \neq f(x); f(x) \in N_2$  then  $\rho_p(p, x) > 0 \forall p \neq x; x \in N_1$ .*

**REMARK 5.6.** In Example 5.2, note that  $c = f((0, c))$  is not connected to all other vertices of  $N_2$ . Observe that  $(0, c)$  is not connected to all other vertices of  $N_1$ .

**DEFINITION 5.7.** Let  $f : N_1 \rightarrow N_2$  be an onto nearring homomorphism,  $\nu$  be a fuzzy ideal of  $N_1$  and  $t \in (\alpha, \beta]$ . Then  $f$  is said to preserve vertex covers of an equiprime fuzzy graph if for each  $\nu_t$  vertex cover of  $(N_{1_e}, \nu, \rho_e, t)$ ,  $f(\nu)_t$  is a vertex cover of  $(N_{2_e}, f(\nu), f(\rho)_e, t)$ .



REMARK 5.8. In Example 5.2, note that  $\nu_t = \{(0, 0), (0, b)\}$  is a vertex cover of  $(N_{1e}, \nu, \rho_e, t)$  and  $f(\nu)_t = \{0, b\}$  is a vertex cover of  $(N_{2e}, f(\nu), f(\rho)_e, t)$ .

PROPOSITION 5.9. *Let  $f : N_1 \rightarrow N_2$  be a nearring homomorphism,  $\nu$  be an equiprime fuzzy ideal of  $N_1$  and  $t \in (\alpha, \beta]$ . If (i)  $f$  preserves vertex covers, (ii)  $f(\nu)_t$  is equisemiprime, then  $f(\nu)$  is an equiprime fuzzy ideal of  $N_2$ .*

*Proof.* Let  $\nu$  be an equiprime fuzzy ideal of  $N_1$ . Then  $f(\nu)$  is a fuzzy ideal of  $N_2$  with the same thresholds as that of  $\nu$ . Suppose  $\nu_t = N_1$ . Then  $f(\nu)_t = f(\nu)_t = N_2$  is an equiprime ideal of  $N_2$ . Then  $f(\nu)$  is an equiprime fuzzy ideal of  $N_2$ . Now suppose  $\nu_t \subset N_1$ . As  $\nu$  is an equiprime fuzzy ideal of  $N_1$  we get  $\nu_t$  is an equiprime ideal of  $N_1$ . By Proposition 3.16 (i) we get  $\nu_t$  is a vertex cover of  $(N_{1e}, \nu, \rho_e, t)$ . As  $f$  preserves vertex cover we get  $f(\nu)_t$  is a vertex covers of  $(N_{2e}, f(\nu), f(\rho)_e, t)$ . Let  $(p, x) \in E(N_2, f(\nu), f(\rho)_e, t)$ . Then  $p \in f(\nu)_t$  or  $x \in f(\nu)_t$ . Then by Proposition 3.9, we get  $p$  is connected to all other vertices of  $(N_{2e}, f(\nu), f(\rho)_e, t)$  or  $x$  is connected to all other vertices of  $(N_{2e}, f(\nu), f(\rho)_e, t)$ . Hence  $(N_{2e}, f(\nu), f(\rho)_e, t)$  is ideal symmetric. Let  $p \in N_2$  such that  $p$  is connected to all other vertices of  $(N_{2e}, f(\nu), f(\rho)_e, t)$ . Then for all  $r \in N_2$  we get  $prx - pr0 \in f(\nu)_t$  for all  $x$  in  $N_2$ . Suppose  $p = x$ . Then  $p \in f(\nu)_t$  (since  $f(\nu)_t$  is equisemiprime). As  $f$  is onto,  $p = f(p_1)$  for some  $p_1 \in N_1$ . Now  $f(p_1)$  is connected to all other vertices of  $(N_{2e}, f(\nu), f(\rho)_e, t)$ . Then by Proposition 5.5, we get  $p_1$  is connected to all other vertices of  $(N_{1e}, \nu, \rho_e, t)$ . Then for all  $r_1 \in N_1$  we get  $(p_1 r_1 x_1 - p_1 r_1 0_1) \in \nu_t$  for all  $x_1 \in N_1$ . Suppose  $\nu_t = N_1$ . Then  $f(\nu)_t = N_2$  is an equiprime ideal of  $N_2$ . Let  $\nu_t \subset N_1$ . Choose  $x_1 \in N_1 \setminus \nu_t$ , then  $p_1 \in \nu_t$ . (As  $\nu$  is an equiprime fuzzy ideal of  $N_1$ ,  $\nu_t$  is an equiprime ideal of  $N_1$ .) Then  $f(p_1) \in f(\nu)_t$  and  $p \in f(\nu)_t$ . Hence  $(N_{2e}, f(\nu), f(\rho)_e, t)$  satisfies all conditions of the Proposition 3.12. Therefor  $f(\nu)$  is an equiprime fuzzy ideal of  $N_2$ .  $\square$

REMARK 5.10. In Example 5.2, note that  $\nu$  is an equiprime fuzzy ideal of  $N_1$  and  $f(\nu)$  is an equiprime fuzzy ideal of  $N_2$ .

ACKNOWLEDGEMENT. We thank the anonymous reviewers and the editor for their constructive comments and suggestions for improving the paper.

The first author thanks St. Joseph Engineering College, Vamanjoor Mangaluru, for encouragement. The second and third authors thank Manipal Institute of Technology and Manipal Academy of Higher Education for their encouragement.

#### REFERENCES

- [1] M. Akram, *m-polar fuzzy graphs*, Stud. Fuzziness Soft Comput., **371** (2019).
- [2] M. Akram, *On T-fuzzy ideals in nearrings*, Int. J. Math. Math. Sci., **2007** (2007), Article ID 73514.
- [3] M. Akram, W. A. Dudek, *Interval valued fuzzy graphs*, Comput. Math. Appl., **61** (2011), 289–299.
- [4] M. Akram, S. Nawaz, *On soft fuzzy graphs*, Ital. J. Pure Appl. Math., **34** (2015), 497–514.
- [5] F. W. Anderson, K. R. Fuller, *Rings and Categories of Modules*, Second edition, Springer-Verlag, USA, 1992.
- [6] I. Beck, *Coloring of commutative rings*, J. Algebra, **116** (1998), 208–226.

- [7] S. K. Bhakat, P. Das, *Fuzzy subrings and ideals redefined*, Fuzzy Sets Syst., **81** (1996), 383–393.
- [8] S. Bhavanari, S. P. Kuncham, B. S. Kedukodi, *Graph of a nearring with respect to an ideal*, Commun. Algebra, **38** (2010), 1957–1962.
- [9] B. Davvaz, *Fuzzy ideals of nearring with interval valued membership functions*, J. Sci. Islam. Repub. Iran, **12** (2001), 171–175.
- [10] B. Davvaz,  $(\epsilon, \epsilon \vee q)$ -fuzzy subnear-rings and ideals, Soft Comput., **10** (2006), 206–211.
- [11] J. A. Goguen, *L-fuzzy sets*, J. Math. Anal. Appl., (1967), 145–174.
- [12] N. J. Groenewald, *Different prime ideals in near-rings*, Comm. Algebra, **19** (1991), 2667–2675.
- [13] F. Harary, *Graph theory*, Naraosa Publishing House, India 2001.
- [14] P. Hell, J. Nešetřil, *Graphs and Homomorphisms*, Oxford University press New York 2004.
- [15] B. Jagadeesha, B. S. Kedukodi, S. P. Kuncham, *Interval Valued L-fuzzy Ideals based on t-norms and t-conorms*, J. Intell. Fuzzy Systems, **28(6)** (2015), 2631–2641.
- [16] B. S. Kedukodi, B. Jagadeesha, S. P. Kuncham, S. Juglal, *Different prime graphs of a nearring with respect to an ideal*, Nearings, Nearfields and related topics, (2017) 185–203.
- [17] B. S. Kedukodi, S. P. Kuncham, S. Bhavanari, *Nearring ideals, graphs and cliques*, Int. Math. Forum, (2013), 73–83.
- [18] B. S. Kedukodi, S. P. Kuncham, S. Bhavanari, *Equiprime, 3-prime and c-prime fuzzy ideals of nearrings*, Soft Comput., **13** (2009), 933–944.
- [19] B. S. Kedukodi, S. P. Kuncham, B. Jagadeesha, *Interval valued L-fuzzy prime ideals, triangular norms and partially ordered groups*, Soft Comput., **23** (2019), 907–920.
- [20] L. T. Koczy, *Fuzzy graphs in the evaluation and optimization of networks*, Fuzzy Sets Syst., **26** (1992), 307–319.
- [21] J. N. Mordeson, P. S. Nair, *Fuzzy Graphs and Fuzzy Hypergraphs*, Physica-Verlag, Heidelberg, 2000.
- [22] J. N. Mordeson, C. Peng, *Operations on fuzzy graphs*, Inform Sci., **79** (1994), 159–170.
- [23] C. Pang, R. Zhang, Q. Zhang, J. Wang, *Dominating sets in directed graphs*, Inform Sci., **180** (2010), 3647–3652.
- [24] G. Pilz, *Near-Rings*, Revised edition, North Holland, 1983.
- [25] S. P. Redmond, *An ideal-based zero-divisor graph of a commutative ring*, Comm. Algebra, **31** (2003), 4425–4443.
- [26] K. P. Shum, M. Akram, *Intuitionistic fuzzy  $(T,S)$ -ideals in nearrings*, J. Algebra Discrete Struct., **6** (2008), 37–52.
- [27] S. Veldsman, *On equiprime near-rings*, Comm. Algebra, **20(9)** (1992), 2569–2587.

(received 18.11.2021; in revised form 12.09.2022; available online 17.04.2023)

Department of Mathematics, St Joseph Engineering College, Vamanjoor Mangaluru 575028, Karnataka India

*E-mail:* jagadeeshab@sjec.ac.in

Department of Mathematics, Manipal Institute of Technology, MAHE, Manipal, Karnataka 576104, India

*E-mail:* babushrisrinivas.k@manipal.edu

Department of Mathematics, Manipal Institute of Technology, MAHE, Manipal, Karnataka 576104, India

*E-mail:* syamprasad.k@manipal.edu