

ON SPLITTING RINGS FOR AZUMAYA SKEW GROUP RINGS

George Szeto and Lianyong Xue

Abstract. Let B be a ring with 1, G an automorphism group of B of order n for some integer n , $B * G$ the skew group ring over B with a free basis $\{g \mid g \in G\}$, B^G the set of elements in B fixed under G , and \overline{G} the inner automorphism group of $B * G$ induced by G . It is shown that when the center C of B is a G -Galois algebra over C^G with Galois group $G|_C \cong G$ or B is a G -Galois extension of B^G and $n^{-1} \in B$, then, $B * G$ is an Azumaya algebra if and only if so is $(B * G)^{\overline{G}}$, and some splitting rings of $B * G$, $(B * G)^{\overline{G}}$ and B are shown to be the same.

1. Introduction

Let B be a ring with 1, C the center of B , G an automorphism group of B of order n for some integer n , $B * G$ a skew group ring over B with a free basis $\{g \mid g \in G\}$, B^G the set of elements in B fixed under G , \overline{G} the inner automorphism group of $B * G$ induced by G , that is, $\overline{g}(f) = gfg^{-1}$ for each $f \in B * G$ and $g \in G$. We note that \overline{G} restricted to B is G .

In [1] and [2], the Azumaya skew group ring $B * G$ over C^G was characterized in terms of Azumaya Galois extension B of B^G and the H -separable extension $B * G$ of B respectively. Also in [3], the commutator subring of B in $B * G$ was studied. In the present paper, under a Galois condition on B , the Azumaya skew group ring $B * G$ is characterized in terms of the Azumaya fixed subring $(B * G)^{\overline{G}}$ under \overline{G} and the Azumaya coefficient ring B , that is, when C is a G -Galois algebra over C^G with Galois group $G|_C \cong G$ or B is a G -Galois extension of B^G and $n^{-1} \in B$, then, $B * G$ is an Azumaya algebra if and only if so is $(B * G)^{\overline{G}}$.

Let A be an Azumaya algebra. It is well known that any separable maximal commutative subalgebra of A is a splitting ring for A ([4], Theorem 5.5, p. 64). In this paper, we call F a splitting ring for A if F is a separable maximal commutative subalgebra of A . We then show that when C is a G -Galois algebra over C^G with Galois group $G|_C \cong G$, F is a splitting ring for the Azumaya algebra $B * G$ containing C if and only if F is a splitting ring for the Azumaya algebra B . Moreover,

AMS Subject Classification: 16S30, 16W20

Keywords and phrases: Skew group rings, Azumaya algebras, Galois extensions, splitting rings

when B is a G -Galois extension of B^G and $n^{-1} \in B$, F is a splitting ring for the Azumaya algebra $B * G$ containing the center of $(B * G)^{\overline{G}}$, then, F is a splitting ring for $(B * G)^{\overline{G}}$ if and only if G is Abelian. At the end, two examples are constructed to demonstrate the results.

This paper was written under the support of a Caterpillar Fellowship at Bradley University. We would like to thank Caterpillar Inc. for the support.

2. Basic definitions and notations

Throughout this paper, B will represent a ring with 1, G an automorphism group of B , C the center of B , $B * G$ a skew ring in which the multiplication is given by $gb = g(b)g$ for $b \in B$ and $g \in G$, B^G the set of elements in B fixed under G , Z the center of $B * G$, \overline{G} the inner automorphism group of $B * G$ induced by G , that is, $\overline{g}(f) = gfg^{-1}$ for each $f \in B * G$ and $g \in G$. We note that \overline{G} restricted to B is G .

Let A be a subring of a ring B with the same identity 1. We denote $V_B(A)$ the commutator subring of A in B . We call B a separable extension of A if there exist $\{a_i, b_i$ in B , $i = 1, 2, \dots, m$ for some integer $m\}$ such that $\sum a_i b_i = 1$, and $\sum b a_i \otimes b_i = \sum a_i \otimes b_i b$ for all b in B where \otimes is over A , and a ring B is called a H -separable extension of A if $B \otimes_A B$ is isomorphic to a direct summand of a finite direct sum of B as a B -bimodule. An Azumaya algebra is a separable extension of its center. B is called a G -Galois extension of B^G if there exist elements $\{c_i, d_i$ in B , $i = 1, 2, \dots, m\}$ for some integer m such that $\sum_{i=1}^m c_i g(d_i) = \delta_{1,g}$. The set $\{c_i, d_i\}$ is called a G -Galois system for B . B is called a DeMeyer-Kanzaki G -Galois extension if B is an Azumaya C -algebra and C is G -Galois algebra with $G|_C \cong G$. If A is an Azumaya C -algebra and S is a commutative C -algebra such that $A \otimes_C S \cong \text{Hom}_S(E, E)$ for some S -progenerator E , then S is called a splitting ring for the Azumaya algebra A . It is well known (Theorem 5.5 in [4] on p. 64) that any separable maximal commutative subalgebra of A is a splitting ring for A . In the present paper, S is called a splitting ring for A if S is a separable maximal commutative subalgebra of A .

3. Characterizations of Azumaya skew group rings

In this section we shall characterize an Azumaya skew group ring $B * G$ in terms of $(B * G)^{\overline{G}}$ and B under a Galois condition that C is a G -Galois algebra over C^G with Galois group $G|_C \cong G$ or B is a G -Galois extension of B^G and $n^{-1} \in B$. We begin with a Lemma.

LEMMA 3.1. *If C is a G -Galois algebra over C^G with Galois group $G|_C \cong G$, then*

- (a) $B * G$ is H -separable over B .
- (b) $B * G$ is H -separable over $(B * G)^{\overline{G}}$.
- (c) The center of $B * G$, $Z = C^G$.
- (d) $V_{B * G}(C) = B$.

Proof. (a) Since C is a G -Galois algebra over C^G with Galois group $G|_C \cong G$ and $C \subseteq V_{B * G}(B)$, $V_{B * G}(B)$ is \overline{G} -Galois extension of $(V_{B * G}(B))^{\overline{G}}$ with the same Galois system as C . Hence, $B * G$ is H -separable extension of B by ([3], Theorem 1).

(b) Since C is a G -Galois extension of C^G with Galois group $G|_C \cong G$, $B * G$ is a \overline{G} -Galois extension of $(B * G)^{\overline{G}}$ with the same Galois system as C . But \overline{G} acts on $B * G$ is inner, so $B * G$ is H -separable extension of $(B * G)^{\overline{G}}$ by ([7], Corollary 3).

(c) By (a), $B * G$ is H -separable over B . Moreover, B is a direct summand of $B * G$ as a left B -module, so B satisfies the double centralizer property in $B * G$ ([8], Proposition 1.2), that is, $B = V_{B * G}(V_{B * G}(B))$. This implies that the center of $B * G$ is contained in B . Thus, $Z = C^G$.

(d) Clearly, $B \subseteq V_{B * G}(C)$. Conversely, for each $\sum_{g \in G} b_g g$ in $V_{B * G}(C)$, we have $c(\sum_{g \in G} b_g g) = (\sum_{g \in G} b_g g)c$ for each c in C , so $cb_g = b_g g(c)$, that is, $b_g(c - g(c)) = 0$ for each $g \in G$ and $c \in C$. But C is a commutative G -Galois extension of C^G , so the ideal of C generated by $\{c - g(c) \mid c \in C\}$ is C ([4], Proposition 1.2-(5)). Thus $b_g = 0$ for each $g \neq 1$. But then $\sum_{g \in G} b_g g = b_1 \in B$. Hence $V_{B * G}(C) \subseteq B$, and so $V_{B * G}(C) = B$. ■

THEOREM 3.2. *Assume C is a G -Galois algebra over C^G with Galois group $G|_C \cong G$. The the following statements are equivalent:*

- (1) $B * G$ is Azumaya.
- (2) $(B * G)^{\overline{G}}$ is Azumaya.
- (3) B is Azumaya.

Proof. (1) \iff (2). Since C is a G -Galois algebra over C^G with Galois group $G|_C \cong G$, there exists an element $c \in C$ such that $\text{Tr}_G(c) = 1$, where $\text{Tr}_G(\)$ is the trace of G ([4], Corollary 1.3-(1)). By Lemma 3.1-(b), $B * G$ is H -separable over $(B * G)^{\overline{G}}$ and $B * G$ is a finitely generated and projective left module over $(B * G)^{\overline{G}}$ ([5], Theorem 1), so (2) \implies (1) by ([6], Theorem 1). Conversely, since the restriction of \overline{G} to C is G , $(B * G)^{\overline{G}}$ is a direct summand of $B * G$ as a $(B * G)^{\overline{G}}$ -bimodule by using the fact that $\text{Tr}_G(c) = 1$. Thus the separability of $B * G$ over Z implies the separability of $(B * G)^{\overline{G}}$ over Z by the argument as given on p. 120 in [5]. Since Z is contained in the center of $(B * G)^{\overline{G}}$, $(B * G)^{\overline{G}}$ is Azumaya. This proves (1) \implies (2).

(1) \implies (3). Assume $B * G$ is Azumaya. Since C is a G -Galois algebra over C^G , $Z = C^G$ by Lemma 3.1-(c). Hence $B * G$ is an Azumaya C^G -algebra. By Lemma 3.1-(d), $V_{B * G}(C) = B$. Therefore, B is a separable C^G -algebra (for C is a separable C^G -algebra) by the commutator theorem for Azumaya algebras ([4], Theorem 4.3, p. 57). Thus B is an Azumaya algebra.

(3) \implies (1). Since C is a $G|_C$ -Galois algebra over C^G , $B * G$ is a H -separable extension of B by Lemma 3.1-(a). By hypothesis, B is an Azumaya C -algebra, so $B * G$ is a separable extension over C by the transitivity of separable extensions. Noting that C is a separable C^G -algebra (for it is G -Galois), we conclude that $B * G$

is a separable extension of C^G . Moreover, by Lemma 1-(c), $Z = C^G$, so $B * G$ is an Azumaya C^G -algebra. ■

THEOREM 3.3. *Let B be a G -Galois extension of B^G and $n^{-1} \in B$. Then, $B * G$ is an Azumaya algebra if and only if so is $(B * G)^{\overline{G}}$. In this case, the center of $(B * G)^{\overline{G}}$ is the center of ZG where Z is the center of $B * G$.*

Proof. Since $n^{-1} \in B$, $\text{Tr}_G(n^{-1}) = 1$. By hypothesis B is a G -Galois extension of B^G , so $B * G$ is a \overline{G} -Galois extension of $(B * G)^{\overline{G}}$ with an inner Galois group \overline{G} with the same Galois system as B . Thus the argument in the proof of (1) \iff (2) in Theorem 3.2 implies that $B * G$ is an Azumaya algebra if and only if so is $(B * G)^{\overline{G}}$.

Next, we calculate the center of $(B * G)^{\overline{G}}$. Let Z be the center of $B * G$. Then the center of $(B * G)^{\overline{G}} = V_{(B * G)^{\overline{G}}}((B * G)^{\overline{G}}) = (B * G)^{\overline{G}} \cap V_{B * G}((B * G)^{\overline{G}}) = (B * G)^{\overline{G}} \cap V_{B * G}(V_{B * G}(ZG))$. Since $n^{-1} \in B$, ZG is a separable Z -algebra. Hence $V_{B * G}(V_{B * G}(ZG)) = ZG$ because $B * G$ is an Azumaya Z -algebra ([4], Theorem 4.3, p. 57). Thus, the center of $(B * G)^{\overline{G}} = (B * G)^{\overline{G}} \cap (ZG) = V_{B * G}(ZG) \cap (ZG) = V_{ZG}(ZG) =$ the center of ZG . ■

4. Splitting rings

In this section, we shall show that some splitting rings for $B * G$, $(B * G)^{\overline{G}}$ and B are the same. Recall that a splitting ring is a separable maximal commutative subalgebra. We first give a result on the splitting rings for any Azumaya algebra.

THEOREM 4.1. *Let A be an Azumaya C -algebra and D a separable commutative subalgebra of A . Then (i) $V_A(D)$ is an Azumaya D -algebra, and (ii) F is a splitting ring for A containing D if and only if F is a splitting ring for $V_A(D)$ over D .*

Proof. (i) Since A is an Azumaya C -algebra and D a separable subalgebra of A , $V_A(V_A(D)) = D$ and $V_A(D)$ is separable subalgebra of A by the commutator theorem for Azumaya algebras ([4], Theorem 4.3, p. 57). Since D is a commutative subalgebra of A , $C \subset D \subset$ the center of $V_A(D)$. Hence $V_A(D)$ is separable over D . Moreover, the center of $V_A(D) = V_{V_A(D)}(V_A(D)) \subset V_A(V_A(D)) = D$; and so the center of $V_A(D) = D$, that is, $V_A(D)$ is an Azumaya D -algebra.

(ii) (\implies) Let F be a splitting ring for A containing D . Then $D \subset F$ and $F = V_A(F)$, and so $F = V_A(F) \subset V_A(D)$. Hence $V_{V_A(D)}(F) = V_A(D) \cap V_A(F) = V_A(F) = F$. Thus F is a maximal commutative subalgebra of $V_A(D)$. Moreover, since F is separable over C and $C \subset$ the center of $V_A(D) = D \subset F =$ the center of F , F is separable over D . Thus, F is splitting ring for $V_A(D)$ over D .

(\impliedby) Let F be splitting ring for $V_A(D)$ over D . Then $D \subset F$ and $F = V_{V_A(D)}(F)$, and so $V_A(F) \subset V_A(D)$. Hence $V_A(F) = V_A(D) \cap V_A(F) = V_{V_A(D)}(F) = F$. Thus F is a maximal commutative subalgebra of A . Moreover, since F is separable over D and D is separable over C , F is separable over C . Therefore, F is splitting ring for A . ■

THEOREM 4.2. *Assume B is a DeMeyer-Kanzaki G -Galois extension (that is, B is an Azumaya C -algebra and C is a G -Galois extension of C^G with $G|_C \cong G$). Then, F is a splitting ring for the Azumaya algebra $B * G$ containing C if and only if F is a splitting ring for the Azumaya algebra B .*

Proof. (\implies) Assume F is a splitting ring for the Azumaya algebra $B * G$ containing C . Then $C \subseteq F$ and $F = V_{B * G}(F)$. Hence $F = V_{B * G}(F) \subseteq V_{B * G}(G)$. Since C is a G -Galois extension of C^G , $V_{B * G}(C) = B$ by Lemma 3.2-(d). Thus $V_{B * G}(F) \subseteq V_{B * G}(C) = B$. Therefore $V_{B * G}(F) = V_B(F)$. But then $F = V_{B * G}(F) = V_B(F)$; and so F is a splitting ring for B .

(\impliedby) Let F be a splitting ring for the Azumaya algebra B . Then $C \subseteq F$ and $F = V_B(F)$. Hence $V_{B * G}(F) \subseteq V_{B * G}(C)$. By Lemma 3.2-(d) again, $V_{B * G}(C) = B$, so $V_{B * G}(F) \subseteq V_{B * G}(C) = B$. Thus $V_{B * G}(F) = V_B(F)$; and so $F = V_B(F) = V_{B * G}(F)$. Therefore, F is a splitting ring for the Azumaya algebra $B * G$ containing C . ■

Next, we consider another Galois condition on B .

THEOREM 4.3. *Let B be a G -Galois extension of B^G , $n^{-1} \in B$ and $B * G$ an Azumaya algebra. Then, F is a splitting ring for $B * G$ containing D , where D is the center of $(B * G)^{\overline{G}}$ if and only if F is a splitting ring for $V_{B * G}(D)$.*

Proof. This is an immediate consequence of Theorem 4.1-(ii) for the Azumaya algebra $B * G$. ■

COROLLARY 4.4. *Assume B is a G -Galois extension of B^G , $n^{-1} \in B$ and $B * G$ an Azumaya algebra. Let G be an Abelian group. Then, F is a splitting ring for $B * G$ containing ZG if and only if F is a splitting ring for $(B * G)^{\overline{G}}$.*

Proof. Since G is Abelian, $n^{-1} \in B$ and Z is the center of $B * G$, ZG is a commutative separable subalgebra. Let $D = ZG$. Then D is the center of $(B * G)^{\overline{G}}$ by Theorem 3.4. Moreover, $V_{B * G}(D) = V_{B * G}(ZG) = (B * G)^{\overline{G}}$, so by Theorem 4.3, F is a splitting ring for $B * G$ containing $ZG (= D)$ if and only if F is a splitting ring for $(B * G)^{\overline{G}} (= V_{B * G}(D))$. ■

THEOREM 4.5. *Assume B is a G -Galois extension of B^G , $n^{-1} \in B$ and $B * G$ is Azumaya algebra. Let F be a splitting ring for $B * G$ containing D , where D is the center of $(B * G)^{\overline{G}}$. Then, F is a splitting ring for $(B * G)^{\overline{G}}$ if and only if G is Abelian.*

Proof. (\implies) Since F is a splitting ring for $B * G$, $F = V_{B * G}(F)$. Now, $F = V_{(B * G)^{\overline{G}}}(F)$, so $F = V_{(B * G)^{\overline{G}}}(F) = (B * G)^{\overline{G}} \cap V_{B * G}(F) = (B * G)^{\overline{G}} \cap F$. Thus $F \subset (B * G)^{\overline{G}}$, and so $F \subset V_{B * G}(ZG)$. Therefore, $V_{B * G}(V_{B * G}(ZG)) \subset V_{B * G}(F) = F$. Since $n^{-1} \in B$, ZG is a separable Z -algebra. Hence $V_{B * G}(V_{B * G}(ZG)) = ZG$ because $B * G$ is an Azumaya Z -algebra ([4], Theorem 4.3, p. 57). Thus, $ZG \subset F$. But F is commutative, so G is Abelian.

(\impliedby) Assume G is Abelian. Since Z is the center of $B * G$, ZG is commutative. Hence $ZG \subset F$, and so $F = V_{B * G}(F) \subset V_{B * G}(ZG)$. Thus $F = V_{B * G}(F) =$

$V_{B * G}(ZG) \cap V_{B * G}(F) = (B * G)^{\overline{G}} \cap V_{B * G}(F) = V_{(B * G)^{\overline{G}}}(F)$. Therefore, F is a splitting ring for $(B * G)^{\overline{G}}$. ■

By Corollary 4.4 and Theorem 4.5, under the hypothesis of Theorem 4.3, two of the following statements imply the third:

- (1) F is a splitting ring for $B * G$ containing the center of $(B * G)^{\overline{G}}$.
- (2) F is a splitting ring for $(B * G)^{\overline{G}}$.
- (3) G is Abelian.

We conclude the present paper with two examples of skew group rings $B * G$ to show the relationship of the splitting rings between $B * G$, B and $(B * G)^{\overline{G}}$.

EXAMPLE 1. Let $B = Q[i, j, k] = Q + Qi + Qj + Qk$ be the quaternion algebra over the rational field Q , $G = \{g_1 = 1, g_i, g_j, g_k \mid g_i(x) = xixi^{-1}, g_j(x) = jxj^{-1}, g_k(x) = kxk^{-1} \text{ for all } x \in B\}$, and $A = B * G$. Then

- (1) B is a G -Galois extension of B^G with G -Galois system $\{\frac{1}{2}, -\frac{1}{2}i, -\frac{1}{2}j, -\frac{1}{2}k; \frac{1}{2}, \frac{1}{2}i, \frac{1}{2}j, \frac{1}{2}k\}$ and $4^{-1} \in B$.
- (2) $B^G = Q$, so A is an Azumaya Q -algebra ([1], Theorem 3.1).
- (3) $D = Q[i] = Q + Qi$ is a commutative separable Q -subalgebra of A .
- (4) $V_A(D) = D + Dg_i + (Qj + Qk)g_j + (Qj + Qk)g_k$ is an Azumaya D -algebra by Theorem 4.1-(i).
- (5) $F = D + Dg_i$ is a splitting ring for $V_A(D)$, so, by Theorem 4.1-(ii), $F = D + Dg_i$ is also a splitting ring for A .
- (6) $(B * G)^{\overline{G}} = V_{B * G}(QG) = QG$ which is a commutative separable subalgebra, so QG is a splitting ring for $(B * G)^{\overline{G}} (= QG)$ and for $B * G$ by Theorem 4.3 (or Corollary 4.4 for G is Abelian).

EXAMPLE 2. Let $M_2(Q)$ be the matrix ring of order 2 over the rational field Q , $B = M_2(Q) \oplus M_2(Q)$, $g: B \rightarrow B$ by $g(a, b) = (b, a)$ for all $(a, b) \in B$. Then,

- (1) g is an automorphism of B of order 2.
- (2) Let $G = \{1, g\}$. Then B is a G -Galois extension of B^G with the Galois system $\{a_1 = (I, 0), a_2 = (0, I); b_1 = (I, 0), b_2 = (0, I)\}$, that is, $a_1b_1 + a_2b_2 = (I, I)$ and $a_1g(b_1) + a_2g(b_2) = (0, 0)$, where I is the identity of $M_2(Q)$ and 0 is the zero matrix in $M_2(Q)$.
- (3) Let C be the center of B . Then $C = Q \oplus Q$, and C is a G -Galois extension of C^G with the same Galois system as B and $C|_G \cong G$.
- (4) $B * G$ is an Azumaya C^G -algebra where $C^G = \{(a, a) \mid a \in Q\}$ since B is an Azumaya C -algebra by Theorem 3.2.
- (5) $(B * G)^{\overline{G}} = C^G + C^G g$.
- (6) Since C is a commutative separable subalgebra of $B * G$, $V_{B * G}(C)$ is an Azumaya C -algebra by Theorem 4.1-(i).
- (7) $V_{B * G}(C) = B$ by Lemma 3.1-(d).

(8) Let $F = Q \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + Q \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$. Then F is a separable maximal commutative subalgebra of $M_2(Q)$, and so $F \oplus F$ is a separable maximal commutative subalgebra of B , that is, $F \oplus F$ is a splitting ring for B . Thus, $F \oplus F$ is a splitting ring for $B * G$ by Theorem 4.2.

REFERENCES

- [1] Alfaro, R. and Szeto, G., *On Galois extensions of an Azumaya algebra*, Comm. in Algebra **25** (6) (1997), 1873–1882.
- [2] Alfaro, R. and Szeto, G., *Skew group rings which are Azumaya*, Comm. in Algebra **23** (6) (1995), 2255–2261.
- [3] Alfaro, R. and Szeto, G., *The centralizer on H -separable skew group rings*, Rings, Extension and Cohomology, Vol. 159, 1995.
- [4] De Meyer, F. R. and Ingraham, E., *Separable Algebras over Commutative Rings*, Vol. 181, Springer Verlag, Berlin, Heidelberg, New York, 1971.
- [5] De Meyer, F. R., *Some notes on the general Galois theory of rings*, Osaka J. Math **2** (1965), 117–127.
- [6] Okamoto, H., *On projective H -separable extensions of Azumaya algebras*, Results in Mathematics **14** (1988), 330–332.
- [7] Sugano, K., *On a special type of Galois extensions*, Hokkaido J. Math. **9** (1980), 123–128.
- [8] Sugano, K., *Note on semisimple extensions and separable extensions*, Osaka J. Math. **4** (1967), 265–270.

(received 08.11.1999.)

Department of Mathematics, Bradley University, Peoria, Illinois 61625, USA