

A NOTE ON IA -AUTOMORPHISMS OF A FINITE p -GROUP

Rasoul Soleimani

Abstract. Let G be a finite group. An automorphism α of G is called an IA -automorphism if $x^{-1}x^\alpha \in G'$ for all $x \in G$. The set of all IA -automorphisms of G is denoted by $\text{Aut}^{G'}(G)$. A group G is called semicomplete if and only if $\text{Aut}^{G'}(G) = \text{Inn}(G)$. In this paper, we obtain certain results on a finite p -group to be semicomplete.

1. Introduction

Let G be a finite group and N a characteristic subgroup of G . Let α be an automorphism of G . If $Ng^\alpha = Ng$ for all g in G , we shall say that α centralizes G/N . We let $\text{Aut}^N(G) = \text{Aut}(G, N)$ denote the centralizer in $\text{Aut}(G)$ of G/N . Clearly $\text{Aut}^N(G)$ is a normal subgroup of $\text{Aut}(G)$, the automorphism group of G , and $\alpha \in \text{Aut}^N(G)$ if and only if $x^{-1}x^\alpha \in N$ for all $x \in G$. The group $\text{Aut}^{G'}(G)$ have been studied by several authors, where G' stands for the derived subgroup of G , see for example [3, 5, 6, 9, 10, 15–17]. Now let M be a normal subgroup of G . We let $\text{Aut}_M(G)$ denote the group of all automorphisms of G centralizing M . Moreover, $\text{Aut}_M^N(G) = \text{Aut}_M(G, N) = \text{Aut}^N(G) \cap \text{Aut}_M(G)$. It is well-known that if G is a finite p -group, then so is the group $\text{Aut}^{G'}(G)$.

In this paper, we study closely the group $\text{Aut}^{G'}(G)$ for a finite p -group G . In Section 2 we give some basic results that are needed for the main results of the paper. In Sections 3 and 4 we prove the main results of the paper and give necessary and sufficient condition for a finite p -group G to be semicomplete when $(G, Z(G))$ is a Camina pair and G' is cyclic.

Throughout the paper all groups are assumed to be finite groups. We use standard notation in group theory. In particular, we use the notation $\text{Hom}(G, A)$ to denote the group of homomorphisms of G into an abelian group A . A group G of order p^m is said to be of maximal class if $m > 2$ and the nilpotency class of G is $m - 1$. A p -group G is said to be extraspecial if $G' = Z(G) = \Phi(G)$ is of order p . Also, a non-abelian group

2010 Mathematics Subject Classification: 20D45, 20D15

Keywords and phrases: Automorphism group; p -group; semicomplete group.

that has no non-trivial abelian direct factor is said to be purely non-abelian. Recall that a group G is called an (internal) central product of its subgroups G_1, \dots, G_n if $G = G_1 \dots G_n$ and $[G_i, G_j] = 1$ for all $1 \leq i < j \leq n$. In this situation, we shall write $G = G_1 * \dots * G_n$. The terms of the lower central series and the upper central series of a group G are respectively denoted as $\Gamma_i(G)$ and $Z_i(G)$. If α is an automorphism of G and x is an element of G , we write x^α for the image of x under α . For a finite group G , $\Omega_i(G)$, $d(G)$, $\mathcal{M}(G)$, $\exp(G)$ and $\text{cl}(G)$ respectively denote the subgroup of G generated by its elements of order dividing p^i , minimal number of generators, the set of all maximal subgroups, the exponent and the nilpotency class of G . Also the size of a finite group G is shown by $|G|$, $o(x)$ for the order of $x \in G$, C_n is the cyclic group of order n and X_{p^3} for non-abelian p -group of order p^3 and exponent p , where p is an odd prime. For $s \geq 1$, we use the notation G^{*s} for the iterated central product defined by $G^{*s} = G * G^{*(s-1)}$ with $G^{*1} = G$, where G is a finite p -group. We also make the convention $G^{*0} = 1$.

2. Some basic results

In this section, we give some known results which will be used in the rest of the paper. An automorphism α of a group G is called central if $x^{-1}x^\alpha \in Z(G)$ for all $x \in G$. The set of all central automorphisms of G is denoted by $\text{Aut}^Z(G)$, where $Z = Z(G)$. The following well-known results will be later used in the paper.

THEOREM 2.1. (*[2, Theorem 1]*) *For a finite purely non-abelian group G , there is a 1-1 correspondence between $\text{Hom}(G, Z(G))$ and $\text{Aut}^Z(G)$, whence $|\text{Hom}(G/G', Z(G))| = |\text{Aut}^Z(G)|$.*

LEMMA 2.2. (*[1, Lemma 2.1]*) *Let G be a finite group and N be a normal subgroup of G such that G/N is abelian. Let $G/N = \langle x_1N \rangle \times \dots \times \langle x_dN \rangle$, where $x_1, \dots, x_d \in G$ and $d = d(G/N)$. If $u_1, \dots, u_d \in Z(N)$ such that*

$$\begin{cases} (x_i u_i)^{n_i} = x_i^{n_i} & 1 \leq i \leq d \\ [x_i, u_j] = [x_j, u_i] & 1 \leq i < j \leq d \end{cases}$$

where $n_i = o(x_i N)$, then the mapping $x_i \mapsto x_i u_i, 1 \leq i \leq d$, can be extended to an automorphism of G leaving N elementwise fixed.

LEMMA 2.3. (*[17, Lemma 2.2]*) *Let G be a group and M, N be normal subgroups of G with $N \leq M$ and $C_N(M) \leq Z(G)$. Then $\text{Aut}_M^N(G) \cong \text{Hom}(G/M, C_N(M))$.*

3. Main results

In this section, we study the group $\text{Aut}^{G'}(G)$ for a finite p -group G . For simplicity, we let $\Gamma_i = \Gamma_i(G)$, for all i .

LEMMA 3.1. *Let G be a finite nilpotent group. If $\alpha \in \text{Aut}^{G'}(G)$ and $a \in \Gamma_i$ ($i = 1, \dots$), then $a^\alpha \equiv a \pmod{\Gamma_{i+1}}$.*

Proof. The result is clearly true for $i = 1$. Proceeding by induction on i , and assume the validity of the lemma for some i . Let $a \in \Gamma_{i+1}$. Then a is a product of terms $b = [y, g]$, such that $y \in \Gamma_i$ and $g \in G$. Now

$$\begin{aligned} b^\alpha &= [y^\alpha, g^\alpha] = [yd, gx], & (d \in \Gamma_{i+1}, x \in G') \\ &= [y, gx]^d [d, gx] \equiv [y, gx]^d & (\text{mod } \Gamma_{i+2}) \\ &= ([y, x][y, g]^x)^d = [y, x]^d [y, g]^{xd} \equiv [y, g]^{xd} & (\text{mod } \Gamma_{i+2}) \\ &= [y, g][[y, g], xd] \equiv [y, g] = b & (\text{mod } \Gamma_{i+2}), \end{aligned}$$

and the lemma follows. \square

THEOREM 3.2. *Let G be a finite nilpotent group of class c . Then*

$$(i) \text{Aut}^{G'}(G) = \text{Aut}_{\Gamma_c}^{G'}(G);$$

$$(ii) \text{Aut}^{\Gamma_c}(G) \leq Z(\text{Aut}^{G'}(G));$$

(iii) $\text{Aut}^{G'}(G)/\text{Aut}^{\Gamma_c}(G)$ is isomorphic to the subgroup of automorphisms in $\text{Aut}^{G'/\Gamma_c}(G/\Gamma_c)$.

Proof. (i) Follows from Lemma 3.1.

To prove (ii), take $\alpha \in \text{Aut}^{\Gamma_c}(G)$ and $\beta \in \text{Aut}^{G'}(G)$. Then for $g \in G$, $g^\alpha = gd$ and $g^\beta = gx$, where $d \in \Gamma_c$ and $x \in G'$. Thus $g^{\alpha\beta} = (gd)^\beta = gxd = gdx$ and $g^{\beta\alpha} = (gx)^\alpha = gdx$, by (i) and since $\text{Aut}(G, \Gamma_c) = \text{Aut}_{G'}(G, \Gamma_c)$. Hence $\alpha\beta = \beta\alpha$ and $\alpha \in Z(\text{Aut}^{G'}(G))$.

(iii) Clearly $\alpha \in \text{Aut}^{G'}(G)$ induces an automorphism $\bar{\alpha}$ in $\frac{G}{\Gamma_c}$, defined by $(g\Gamma_c)^{\bar{\alpha}} = g^\alpha\Gamma_c$. It is easy to see that the mapping $\alpha \mapsto \bar{\alpha}$ defines a homomorphism of $\text{Aut}^{G'}(G)$ into $\text{Aut}(\frac{G}{\Gamma_c}, \frac{G'}{\Gamma_c})$. The kernel of this homomorphism is $\text{Aut}^{\Gamma_c}(G)$, for $\bar{\alpha} = \bar{1}$ if and only if $g^{-1}g^\alpha \in \Gamma_c$, for all $g \in G$, which means that $\alpha \in \text{Aut}(G, \Gamma_c)$. \square

THEOREM 3.3. *Let G be a finite nilpotent group. Then $\text{cl}(\text{Aut}^{G'}(G)) = \text{cl}(G) - 1$.*

Proof. Suppose that $\text{cl}(G) = c$. We use induction on c . For $c = 1$, it is clearly true. Assume that the result holds for any finite nilpotent group of nilpotency class less than c . Hence $\text{cl}(\text{Aut}(\frac{G}{\Gamma_c}, \frac{G'}{\Gamma_c})) = \text{cl}(\frac{G}{\Gamma_c}) - 1 \leq \text{cl}(G) - 2$. Since $\text{Inn}(G) \leq \text{Aut}^{G'}(G)$, $\text{cl}(G) - 1 \leq \text{cl}(\text{Aut}^{G'}(G))$. Now by Theorem 3.2 (ii) and (iii) we have $\text{cl}(G) - 2 \leq \text{cl}(\text{Aut}^{G'}(G)) - 1 = \text{cl}(\frac{\text{Aut}^{G'}(G)}{Z(\text{Aut}^{G'}(G))}) \leq \text{cl}(\frac{\text{Aut}^{G'}(G)}{\text{Aut}(G, \Gamma_c)}) \leq \text{cl}(\text{Aut}(\frac{G}{\Gamma_c}, \frac{G'}{\Gamma_c})) \leq \text{cl}(G) - 2$.

Consequently, $\text{cl}(\text{Aut}^{G'}(G)) - 1 = \text{cl}(G) - 2$ and $\text{cl}(\text{Aut}^{G'}(G)) = \text{cl}(G) - 1$. \square

COROLLARY 3.4. *Let G be a finite p -group of order p^n . Then G is of maximal class if and only if $\text{cl}(\text{Aut}^{G'}(G)) = n - 2$.*

THEOREM 3.5. *Let G be a finite p -group of class c . Then $\text{Aut}_Z^{\Gamma_c}(\Gamma_{c-1}) = \text{Inn}(\Gamma_{c-1})$ if and only if Γ_c is cyclic, where $Z = Z(\Gamma_{c-1})$.*

Proof. By Lemma 2.3, $\text{Aut}_Z^{\Gamma_c}(\Gamma_{c-1}) \cong \text{Hom}(\Gamma_{c-1}/Z(\Gamma_{c-1}), \Gamma_c)$. It is sufficient to prove that $\exp(\Gamma_{c-1}/Z(\Gamma_{c-1})) \leq \exp(\Gamma_c)$. Suppose that $\exp(\Gamma_c) = p^n$ and $g \in \Gamma_{c-1}$ such that $o(gZ(\Gamma_{c-1})) = \exp(\Gamma_{c-1}/Z(\Gamma_{c-1}))$. Now $[g^{p^n}, x] = [g, x]^{p^n} = 1$ for all $x \in G$. So $g^{p^n} \in Z(\Gamma_{c-1})$ and the proof is complete. \square

As an application of Theorem 3.5, we get the following corollary which is the same as [17, Proposition 3.2].

COROLLARY 3.6. *Let G be a finite p -group of class 2. Then $\text{Aut}_Z^{G'}(G) = \text{Inn}(G)$ if and only if G' is cyclic, where $Z = Z(G)$.*

THEOREM 3.7. *Let G be a finite p -group of class c . Then $\text{Aut}_{\Gamma_c}^{\Gamma_c}(\Gamma_{c-1}) = \text{Inn}(\Gamma_{c-1})$ if and only if Γ_c is cyclic and $Z(\Gamma_{c-1}) = \Gamma_c \Gamma_{c-1}^{p^n}$, where $|\Gamma_c| = p^n$.*

Proof. Assume that Γ_c is cyclic and of order p^n . By Theorem 3.5, it is sufficient to prove that $\text{Aut}_{\Gamma_c}^{\Gamma_c}(\Gamma_{c-1}) = \text{Aut}_Z^{\Gamma_c}(\Gamma_{c-1})$, where $Z = Z(\Gamma_{c-1})$. Let $\alpha \in \text{Aut}_{\Gamma_c}^{\Gamma_c}(\Gamma_{c-1})$ and $x \in \Gamma_{c-1}$. We may write $(x^{p^n})^\alpha = (xd)^{p^n} = x^{p^n}$ with $d \in \Gamma_c$, which shows that α fixes any element of $Z(\Gamma_{c-1})$, since $Z(\Gamma_{c-1}) = \Gamma_c \Gamma_{c-1}^{p^n}$. Consequently $\text{Aut}_{\Gamma_c}^{\Gamma_c}(\Gamma_{c-1}) = \text{Aut}_Z^{\Gamma_c}(\Gamma_{c-1}) = \text{Inn}(\Gamma_{c-1})$.

Conversely, suppose that $\text{Aut}_{\Gamma_c}^{\Gamma_c}(\Gamma_{c-1}) = \text{Inn}(\Gamma_{c-1})$. By Theorem 3.5, Γ_c is cyclic. Since $\Gamma_c \leq \Gamma_c \Gamma_{c-1}^{p^n} \leq Z(\Gamma_{c-1}) \leq \Gamma_{c-1}$, it follows that

$$\begin{aligned} \text{Inn}(\Gamma_{c-1}) &\cong \text{Hom}(\Gamma_{c-1}/Z(\Gamma_{c-1}), \Gamma_c) \twoheadrightarrow \text{Hom}(\Gamma_{c-1}/(\Gamma_c \Gamma_{c-1}^{p^n}), \Gamma_c) \\ &\twoheadrightarrow \text{Hom}(\Gamma_{c-1}/\Gamma_c, \Gamma_c) \cong \text{Aut}_{\Gamma_c}(\Gamma_{c-1}, \Gamma_c) = \text{Inn}(\Gamma_{c-1}). \end{aligned}$$

Therefore $\text{Hom}(\Gamma_{c-1}/(\Gamma_c \Gamma_{c-1}^{p^n}), \Gamma_c) \cong \text{Inn}(\Gamma_{c-1})$, which gives $|\Gamma_{c-1}/(\Gamma_c \Gamma_{c-1}^{p^n})| = |\Gamma_{c-1}/Z(\Gamma_{c-1})|$. So $Z(\Gamma_{c-1}) = \Gamma_c \Gamma_{c-1}^{p^n}$, as required. \square

S. Singh, D. Gumber, and H. Kalra [15] gave a necessary and sufficient condition on a finite p -group to be semicomplete. Our next corollary, which is a particular case of Theorem 3.7, gives an another interpretation of this result. This corollary is [17, Theorem 3.3].

COROLLARY 3.8. *Let G be a finite p -group of class 2. Then $\text{Aut}^{G'}(G) = \text{Inn}(G)$ if and only if G' is cyclic and $Z(G) = G'G^{p^n}$, where $|G'| = p^n$.*

We now give an alternative proof for [15, Corollary 2.4].

COROLLARY 3.9. *Let G be a 2-generated finite nilpotent group of class 2. Then any IA-automorphism of G is an inner automorphism.*

Proof. Suppose that $G = \langle a, b \rangle$. Then $G' = \langle [a, b]^g | g \in G \rangle = \langle [a, b] \rangle$ and so G' is cyclic. Since G is a nilpotent group, $G = P_1 \times \dots \times P_n$, where P_i is the Sylow p_i -subgroup of G , for $i = 1, \dots, n$. Thus $G' = P'_k$, $\text{Inn}(G) \cong \text{Inn}(P_k)$ and by Lemma 2.3, $\text{Aut}^{G'}(G) \cong \text{Aut}^{P'_1}(P_1) \times \dots \times \text{Aut}^{P'_n}(P_n) = \text{Aut}^{P'_k}(P_k) \cong \text{Hom}(P_k/P'_k, P'_k)$ for some

$1 \leq k \leq n$. Next by [5, Theorem 3.2], $|\text{Aut}^{G'}(G)| = |G'|^2$ and so $|\text{Aut}^{P'_k}(P_k)| = |P'_k|^2$. Now since $P'_k \leq Z(P_k)$, by [14, Lemma 0.4], if $\exp(P_k/Z(P_k)) = p^m = \exp(P'_k)$, then $P_k/Z(P_k)$ has the form $C_{p^m} \times C_{p^m} \times A$ for some (possibly trivial) abelian p -group A . So by Lemma 2.3, $|\text{Aut}_Z^{P'_k}(P_k)| = |\text{Hom}(P_k/Z(P_k), P'_k)| \geq |P'_k|^2$, where $Z = Z(P_k)$. Thus $\text{Aut}^{P'_k}(P_k) = \text{Aut}_Z^{P'_k}(P_k)$, which together with Corollary 3.6 completes the proof. \square

4. Groups G such that $(G, Z(G))$ is a Camina pair

Camina groups were introduced by A.R. Camina in [4] and were studied in past (see for example [11–13]). Let G be a finite group and N be non-trivial proper normal subgroup of G . Then (G, N) is called a *Camina pair* if $xN \subseteq x^G$ for all $x \in G - N$, where x^G denotes the conjugacy class of x in G . It follows that (G, N) is a Camina pair if and only if $N \subseteq [x, G]$ for all $x \in G - N$, where $[x, G] = \{[x, g] | g \in G\}$.

In this section, we give necessary and sufficient condition for a finite p -group G to be semicomplete when $(G, Z(G))$ is a Camina pair and G' is cyclic. We start with some results of I.D. Macdonald.

LEMMA 4.1. ([12, Lemma 2.1]) *Let (G, H) be a Camina pair and G have class c . Then $H = \Gamma_r(G)$ and $H = Z_{c-r+1}(G)$ for some r satisfying $1 < r \leq c$.*

THEOREM 4.2. ([12, Theorem 2.2]) *Let (G, H) be a Camina pair, $H = Z(G)$, and G have class c . Then $Z_r(G)/Z_{r-1}(G)$ has exponent p whenever $1 \leq r \leq c$.*

THEOREM 4.3. *Let G be a finite p -group such that G' is cyclic and $(G, Z(G))$ is a Camina pair. Then $\text{Aut}^{G'}(G) = \text{Inn}(G)$ if and only if G is an extraspecial p -group or G is isomorphic to a central product $A * X_{p^3}^{*s}$, for some $s \geq 0$, p is an odd prime and A is a 2-generator subgroup which is either a metacyclic group or $A = \langle a \rangle \langle b \rangle \langle c \rangle$, $[a, c] = [b, c] = 1$, $[a, b] = cb^{p^k}$, where $k \geq 1$.*

Proof. Let $(G, Z(G))$ be a Camina pair and $\alpha \in \text{Aut}^Z(G)$, where $Z = Z(G)$. Since $Z(G) \leq G'$, $\frac{Z_2(G)}{Z(G)} \cong \text{Aut}^Z(G) \cap \text{Inn}(G) = \text{Aut}^Z(G)$ and so by Theorem 4.2, $\text{Aut}^Z(G)$ is elementary abelian. Now $Z(G) < Z(M)$ and $C_G(M) = Z(M)$, for all $M \in \mathcal{M}(G)$ [7, Remark 2]. Assume that $|G/\Phi(G)| = p^t$ and $|Z(G)| = p^r$. By [18, Theorem 3.1], $d(Z_2(G)/Z(G)) = d(G)$. Since G is purely non-abelian, we have $p^t = |\text{Aut}^Z(G)| = |\text{Hom}(G/G', Z(G))| = p^{rt}$, by Theorem 2.1. Whence $r = 1$ and $Z(G) \cong C_p$. If $G/Z(G)$ be an abelian then by Corollary 3.8, $G' = Z(G) = \Phi(G) \cong C_p$ and hence G is extraspecial. So we may assume that $G/Z(G)$ is not abelian.

We first assume that $p > 2$. Then by the main theorem of [8], we may write $G = A_1 * A_2 * \dots * A_n * B$, where B is an abelian subgroup, A_1, A_2, \dots, A_n are 2-generator subgroups, and the classes of A_2, \dots, A_n are equal to 2. Now $G = A_1 * A_2 * \dots * A_n$, since $B \leq Z(G) \leq \Phi(G)$. Next for $2 \leq i \leq n$, $(A_i, Z(A_i))$ is a Camina pair since $xZ(A_i) = xZ(G) \subseteq x^G = x^{A_1 \dots A_n} = x^{A_i}$, for all $x \in A_i - Z(A_i)$. Thus $A'_i = Z(A_i) = Z(G) \cong C_p$ and A_i is an extraspecial p -group of order p^3 and exponent

p , where $2 \leq i \leq n$. So by the theorem mentioned earlier, it follows that $G \cong A * X_{p^3}^{*s}$, where $s \geq 0$ and A is a 2-generator subgroup which is either a metacyclic group or $A = \langle a \rangle \langle b \rangle \langle c \rangle$, $[a, c] = [b, c] = 1$, $[a, b] = cb^{p^k}$, $k \geq 1$.

Suppose next that $p = 2$. First we show that $Z(M) \leq Z_2(G)$, for all $M \in \mathcal{M}(G)$. Let $M \in \mathcal{M}(G)$, $g \in G \setminus M$ and $x \in Z(M) \setminus Z(G)$. Since $g^2 \in M$, $[x, G] = [x, M \langle g \rangle] = \{[x, g^i] \mid 0 \leq i < 2\}$. By assumption $Z(G) \subseteq [x, G]$ and $|Z(G)| = 2$. Consequently $Z(G) = [x, G]$ and so $x \in Z_2(G)$. Next let $x \in Z_2(G) \setminus Z(G)$. It follows that $M = C_G(x)$ is a maximal subgroup of G , since $|C_G(x)| = |G|/[x, G] = |G|/2$. Let $(Z_2(G) \cap G')/Z(G) = \langle \bar{t} \rangle$ and $M = C_G(t)$, where $t \in Z_2(G) \cap G'$ and $\bar{t} = tZ(G)$. Then $M \in \mathcal{M}(G)$ and if $g \in G \setminus M$, it follows that $[t, g] \in Z(G)$. Hence $(gt)^2 = g^2 t^2 [t, g] = g^2$, since $o(t) = 4$ and $[t, g] = t^2$. Now since $t \in Z(M)$, the map α sending $g \mapsto gt$ and $m \mapsto m$, for all $m \in M$, can be extended to an automorphism of G by Lemma 2.2, which is an automorphism lying in $\text{Aut}^{G'}(G)$. So that α is an inner automorphism of G induced by an element x_M in G . It follows that $x_M \in C_G(M) = Z(M) \leq Z_2(G)$. This means that $t = g^{-1} g^\alpha = [g, x_M] \in Z(G)$, which is impossible.

Conversely, if G is an extraspecial p -group then by Lemma 2.3, $\text{Aut}^{G'}(G) \cong \text{Hom}(G/G', G') \cong \text{Inn}(G)$, and so G is semicomplete. Next let $G \cong A * X_{p^3}^{*s}$, for some $s \geq 0$ and $p > 2$. Then by Theorem 3.2.(i), Lemma 4.1 and [6, Theorem 3], $\text{Aut}^{G'}(G) = \text{Aut}_{\Gamma_c}^{G'}(G) = \text{Aut}_Z^{G'}(G) = \text{Inn}(G)$, which completes the proof. \square

ACKNOWLEDGEMENT. The author is grateful to the referee for his/her valuable suggestions. The paper was revised according to these comments. This research was in part supported by a grant from Payame Noor University.

REFERENCES

- [1] A. Abdollahi, S.M. Ghoraihi, *On noninner 2-automorphisms of finite 2-groups*, Bull. Aust. Math. Soc. **90**(2) (2014), 227–231.
- [2] J.E. Adney, T. Yen, *Automorphisms of a p -group*, Ill. J. Math. **9**(1965), 137–143.
- [3] S. Bachmuth, *Automorphisms of free metabelian groups*, Trans. Amer. Math. Soc. **118** (1965), 93–104.
- [4] A.R. Camina, *Some conditions which almost characterize Frobenius groups*, Israel J. Math. **31** (1978), 153–160.
- [5] A. Caranti, C.M. Scoppola, *Endomorphisms of two-generated metabelian groups that induce the identity modulo the derived subgroup*, Arch. Math. **56** (1991), 218–227.
- [6] Y. Cheng, *On finite p -groups with cyclic commutator subgroup*, Arch. Math. **39** (1982), 295–298.
- [7] M. Deaconescu, G. Silberberg, *Noninner automorphisms of order p of finite p -groups*, J. Algebra, **250** (2002), 283–287.
- [8] A. A. Finogenov, *Finite p -groups with a cyclic commutator subgroup*, Algebra Logic **34**(2) (1995), 125–129.
- [9] T. A. Fournelle, *Torsion in semicomplete nilpotent groups*, Math. Proc. Cambridge Philos. Soc. **94** (1983), 191–202.
- [10] C.K. Gupta, *IA-automorphisms of two generator metabelian groups*, Arch. Math. **37** (1981), 106–112.
- [11] M.L. Lewis, *On p -group Camina pairs*, J. Group Theory, **15** (2012), 469–483.
- [12] I.D. Macdonald, *Some p -groups of Frobenius and extra-special type*, Israel J. Math. **40** (1981), 350–364.

- [13] I.D. Macdonald, *More on p -groups of Frobenius type*, Israel J. Math. **56** (1986), 335–344.
- [14] M. Morigi, *On the minimal number of generators of finite non-abelian p -groups having an abelian automorphism group*, Comm. Algebra, **23** (1995), 2045–2065.
- [15] S. Singh, D. Gumber, H. Kalra, *IA -automorphisms of finitely generated nilpotent groups*, J. Algebra Appl. **13** (2014), 1450027.
- [16] M. Shabani Attar, *Semicomplete finite p -groups*, Algebra Colloq. **18** (Spec 1) (2011), 937–944.
- [17] R. Soleimani, *On some p -subgroups of automorphism group of a finite p -group*, Vietnam J. Math. **36**(1) (2008), 63–69.
- [18] M.K. Yadav, *On automorphisms of finite p -groups*, J. Group Theory, **10** (2007), 859–866.

(received 13.06.2017; in revised form 15.01.2018; available online 02.02.2018)

Department of Mathematics, Payame Noor University (PNU), 19395-3697, Tehran, Iran
E-mail: r_soleimani@pnu.ac.ir, rsoleimani@yahoo.com